

SUBDIVISIONS IN APEX GRAPHS

Elad Aigner-Horev
elad.horev@uni-hamburg.de

Department of Mathematics
Hamburg university

Abstract. The Kelmans-Seymour conjecture states that the 5-connected nonplanar graphs contain a subdivided K_5 . Certain questions of Mader propose a “plan” towards a possible resolution of this conjecture. One part of this plan is to show that a 5-connected nonplanar graph containing K_4^- or $K_{2,3}$ as a subgraph has a subdivided K_5 . Recently, Ma and Yu showed that a 5-connected nonplanar graph containing K_4^- as a subgraph has a subdivided K_5 . We take interest in $K_{2,3}$ and prove that a 5-connected nonplanar apex graph containing $K_{2,3}$ as a subgraph has a subdivided K_5 .

Keywords. (Rooted) Subdivisions, Nonseparating paths, Apex graphs.

PREAMBLE. Whenever possible notation and terminology are that of [1]. Throughout, a graph is always simple, undirected, and finite. G always denotes a graph. A subdivided G is denoted TG and the *branch* vertices of a TG are its vertices at least 3-valent. A TG is said to be *rooted* at a prescribed vertex set X , if the latter coincides with its branch vertices.

§1 Introduction. A refinement of Kuratowski’s theorem postulated by the Kelmans-Seymour conjecture (1975) is that: *the 5-connected nonplanar graphs contain a TK_5* . Certain questions of Mader [4] seem to suggest a two phase plan towards a possible resolution of the Kelmans-Seymour conjecture.

MADER’S “PLAN”.

Phase I. Find a small set of graphs $\mathcal{H}$ such that every graph with at least $\lfloor \frac{5}{2}n \rfloor$ edges contains a TK_5 or a member of $\mathcal{H}$ as a subgraph.

Phase II. Prove: for every $H \in \mathcal{H}$, if G is a 5-connected nonplanar graph with H as a subgraph, then $TK_5 \subseteq G$.

The questions of Mader [4] and the discussion thereafter suggest to examine $\mathcal{H} = \{K_4^-, K_{2,3}\}$, where K_4^- denotes K_4 with a single edge removed. We are not aware of any mathematics that would indicate as to which phase of the above plan one should consider first. Nevertheless, recently, Ma and Yu [2, 3] proved the following.

1.1. (Ma-Yu [2, 3])

A 5-connected nonplanar graph containing K_4^- as a subgraph, contains a TK_5 .

We take interest in $K_{2,3}$ and ask *is it true that a nonplanar 5-connected graph containing a $K_{2,3}$ as a subgraph contains a TK_5* ? Using 1.1, we answer this question in the affirmative for apex graphs, where a graph is apex if it contains a vertex removal of which results in a planar graph.

1.2. *A 5-connected nonplanar apex graph containing $K_{2,3}$ as a subgraph contains a TK_5 .*

The restriction of **1.2** to apex graphs is not artificial. Indeed, the conjecture being open for so many years suggests that it does not stand to reason to consider special cases of it. We contend that apex graphs rise quite naturally; for instance, when trying to determine properties of a minimal counter example to the conjecture.

Our approach to prove **1.2** is structural and as a result most of its significant parts do not use the assumption that the graph is apex. The generalization of the parts of the proof that do rely on this assumption coincide with a well-known problem proposed by Yu [9]. To emphasize this, the proof is written in a modular manner so that the dependency on planarity (i.e., the apex assumption) can be easily singled out. Next, we make this more precise as we outline the highlights of our approach.

I. Essentially, to prove **1.2**, we complete a $K_{2,3}$ into a TK_5 . Such an approach brings forth a problem which we have come to call the *rooted nonseparating path problem*: for integer $k > 0$, $x, y \in V(G)$, and $Y \subset V(G) \setminus \{x, y\}$ (the roots), is there an induced xy -path P in G satisfying $\kappa(G - P) \geq k$ and $V(P) \cap Y = \emptyset$? This problem is at the heart of the Ma-Yu [2, 3] proof for **1.1**; it will also be at the heart of our proof for **1.2**. In particular, we prove **4.7** which we propose to be of independent interest.

II. The assertion of **4.7** allows us to assume that if $K_{2,3} \cong K \subseteq G$, then, essentially, G is comprised of two parts, i.e., $G = H \cup P$, where P is an induced path linking the 3-valent vertices of K meeting no 2-valent vertex of K , and $H = G - P$ is a chain of blocks (i.e., maximal 2-connected components of $G - P$). The following is then one of our main results.

1.3. Suppose G is a 5-connected graph such that $K_{2,3} \cong K \subseteq G$. If G has an induced path P linking the 3-valent vertices of K meeting no 2-valent vertex of K such that $\kappa(G - P) \geq 2$, then $TK_5 \subseteq G$.

Note that **1.3** is not limited to apex nor to nonplanar graphs and in our opinion would serve as a good stepping stone towards generalizing **1.2** for nonapex graphs. The characterization of 2-connected graphs containing no circuit through 3 prescribed elements (edges or vertices) is the main engine in our proof of **1.3** (see §3.1).

III. Sadly, **1.3** will not be sufficient for us as we do not know how to guarantee existence of such a path. When confined to apex graphs a new structural tool is made available; such is the characterization, obtained by Yu [9], of pairs (G, W) such that G is a 4-connected plane graph, $W \subseteq V(G)$, $|W| = 4$, and G has no TK_4 rooted at W (see §4.2 for details).

At this point of the proof, we have two descriptions of the graph G . One description is that which emerges from applying Yu's result in [9] in certain ways; the second is that which emerges from **1.3** and **4.7** that $G = H \cup P$ where H and P are as above and $\kappa(H) = 1$. A comparison between these two descriptions is then drawn as to arrive at a conclusion of our proof.

Appealing to the result of Yu [9] (and the analysis that it allows us to carry) is essentially the sole place of the proof that relies on planarity and the graph being apex. In [9], Yu proposes the problem of generalizing his result to pairs (G, W) as above where G is not necessarily planar.

It is worth to note that using this characterization, Yu [9] proved that every 6-connected apex graph contains a TK_5 . This was then superseded by Mader [4] who proceeded to prove that every 6-connected graph contains a TK_5 .

IV. Finally, **1.1** and a variant of the following recent result of Ma and Yu [2] will be used extensively throughout our arguments for **1.2**. Indeed, these two theorems of Ma and Yu form a useful tool in our analysis.

1.4. (Ma-Yu [2])

Let G be a 5-connected nonplanar graph such that $G = G_1 \cup G_2$, $|V(G_1 \cap G_2)| = 5$. If $|V(G)| > |V(G_2)| \geq 7$ and G_2 has a planar representation with $V(G_1 \cap G_2)$ incident with a common face, then $TK_5 \subseteq G$.

ORGANIZATION. Additional notation, terminology, and required previous results are detailed in §2. A proof of **1.3** is provided in §3. A proof of **1.2** is provided in §4.

§2 Preliminaries. Throughout, whenever a claim regarding plane graphs is applied to a planar graph G , we always assume that the application is done with respect to some embedding of G .

AGREEMENT. We always assume that $G + e = G$ if $e \in E(G)$. In addition to that, if $e \in E(G)$ and we write $G' = G + e$ and then $G'' = G' - e$ in an argument, then the agreement is that $G' = G = G''$. We use this agreement in §3 and §4.4.

SUBGRAPHS. Let H be a subgraph of G , denoted $H \subseteq G$. The boundary of H , denoted by $\text{bnd}_G H$ (or simply by $\text{bnd} H$), is the set of vertices of H incident with $E(G) \setminus E(H)$. By $\text{int}_G H$ (or simply $\text{int} H$) we denote the subgraph induced by $V(H) \setminus \text{bnd} H$. If $v \in V(G)$, then $N_H(v)$ denotes $N_G(v) \cap V(H)$.

Given $U \subseteq V(G) \cup E(G)$, we write $U \cap H$ to denote $U \cap (V(H) \cup E(H))$. We also write $x \in G$ to denote that x is an element of G , i.e., an edge or a vertex.

PATHS AND CIRCUITS. For $X, Y \subseteq V(G)$, an (X, Y) -path is a simple path with one end in X and the other in Y internally-disjoint of $X \cup Y$. If $X = \{x\}$, we write (x, Y) -path. If $|X| = |Y| = k \geq 1$, then a set of k pairwise disjoint (X, Y) -paths is called an (X, Y) - k -linkage. Also, if $x \in V(G)$ and $Y \subseteq V(G) \setminus \{x\}$, then by (x, Y) - k -fan we mean a set of $k \geq 1$ (x, Y) -paths with only x as a common vertex.

The *interior* of an xy -path P is the set $V(P) \setminus \{x, y\}$ and is denoted $\text{int} P$. For $u, v \in V(P)$, we write uPv to denote the vertex set of the uv -subpath of P and is called the uv -segment of P . We write (uPv) to denote $uPv \setminus \{u, v\}$, and in a similar manner the segment (uPv) and the segment uPv are defined.

A uv -segment of P nests in a $u'v'$ -segment of P if $uPv \subseteq u'Pv'$. These segments are said to *overlap* if x, u, u', v, v', y appear in this order along P .

Let $H_1 \subseteq G - P$. A vertex $u \in V(P)$ adjacent to H_1 is called an *attachment vertex* of H_1 on P . If H_1 has ≥ 2 attachments on P , then the attachments v_1, v_2 such that $v_1 P v_2$ is maximal are called the *extremal* attachments of H_1 on P ; the set $v_1 P v_2$ is then called the *extremal segment* of H_1 on P .

We say that H_1 *neests* in $H_2 \subseteq G - P$ with respect to P if the extremal segment of H_1 on P nests in the extremal segment of H_2 on P . These subgraphs are said to *overlap* with respect to P if their extremal segments on P overlap.

Given a set $A \subseteq V(G) \cup E(G)$, we refer to a circuit containing A as an A -*circuit*. For a circuit C embedded in the plane we denote by $\text{int}C$ and $\text{ext}C$ the regions of the plane bounded by C and appearing in its interior and exterior, respectively.

BRIDGES. Let $H \subseteq G$. By H -*bridge* we mean either an edge $uv \notin E(H)$ and $u, v \in V(H)$ or a connected component of $G - H$. In the latter case, the H -bridge is called *nontrivial*. The vertices of H adjacent to an H -bridge B are called the *attachment vertices* of B . A uv -path internally-disjoint of H with $u, v \in V(H)$, is called an H -*ear*.

HAMMOCKS AND DISCONNECTORS. A k -*hammock* in G is a connected subgraph whose boundary consists of $k \geq 1$ vertices; called *ends*. A hammock coinciding with its boundary is called *trivial*. In this sense, a trivial 2-hammock is an edge. For 2-hammocks we also refer to $G - e$ as a trivial 2-hammock, where $e \in E(G)$. Clearly, a graph with no nontrivial 2-hammocks is 3-connected, and so on.

Let G have $\kappa(G) = 2$. Then G is a union of 2-hammocks. An end-to-end path in a 2-hammock will be called a *through path*. Being not necessarily 2-connected, a 2-hammock becomes such after an ear linking its ends is appended. Equivalently,

$$\text{any edge of a 2-hammock is traversed by a through path.} \quad (2.1)$$

If G is connected and $D \subseteq V(G)$ satisfies $|D| = k$ and $G - D$ is disconnected, then D is called a k -*disconnecter*. The boundary of a k -hammock H is a k -disconnecter, unless H is trivial.

PLANAR HAMMOCKS. The argument of Ma and Yu [2] for **1.4** asserts the following.

2.2. *Let G be a 5-connected nonplanar graph such that $G = G_1 \cup G_2$, $|V(G_1 \cap G_2)| = 5$. If $|V(G)| > |V(G_2)| \geq 7$ and G_2 has a an embedding in the closed disc with $V(G_1 \cap G_2)$ appearing on the boundary of the disc, then $TK_5 \subseteq G$.*

Indeed, the argument of Ma and Yu for **1.4** shows that the existence of a separation as in **2.2** can be translated into a TK_5 , where the main engine is [2, Theorem 4.3].

A 5-hammock H of G is called a *planar hammock* if $|V(G)| > |V(H)| \geq 7$ and H has an embedding in the closed disc with $\text{bnd}H$ appearing on the boundary of the disc. For us it will be more convenient to use the following.

2.3. *A 5-connected nonplanar graph containing a planar hammock contains a TK_5 .*

BLOCKS. The *blocks* of G are its maximal 2-connected components. We refer to a single edge as being 2-connected. The boundary of a block of G is the set of cut vertices of G contained in the block.

If G is connected, then the blocks and cut vertices of G define a unique tree called the *block tree* of G ; the leaves of which are blocks. A graph is called a *chain* of blocks if its block

tree is a (simple) path; and is said to be a *claw-chain* of blocks if its block tree has precisely 3 leaves. By *xy-chain* we mean a chain of blocks with each member of $\{x, y\}$ met by a leaf block of the chain. A subgraph of a chain that is also a chain is called a *subchain*.

§3 Proof of 1.3. Suppose G is a graph such that

$$K_{2,3} \cong K \subseteq G, V(K) = \{x_1, x_2, x_3, x_4, x_5\} \text{ such that } d_K(x_1) = d_K(x_2) = 3. \quad (3.1)$$

We observe that if there is an x_1x_2 -path P in $G - \{x_3, x_4, x_5\}$ such that

(F.1) $G - P$ contains an $\{x_3, x_4, x_5\}$ -circuit, then $TK_5 \subseteq G$.

(F.2) Also, if there is a $v \in \text{int}P$ such that $G - P + v$ contains a $Z \cup \{v\}$ -circuit C satisfying $V(C) \cap \{x_3, x_4, x_5\} = Z$ and $|Z| = 2$, then $TK_5 \subseteq G$.

We restate **1.3** as follows.

3.2. Let G be a 5-connected graph with $K \subseteq G$ as in (3.1). If G has an induced x_1x_2 -path P satisfying $x_3, x_4, x_5 \notin V(P)$ and $\kappa(G - P) \geq 2$, then $TK_5 \subseteq G$.

If $TK_5 \not\subseteq G$ and G, P are as in **3.2**, then (F.1) fails in the sense that $G - P$ contains no $\{x_3, x_4, x_5\}$ -circuit. By **3.4**, which is the characterization of 2-connected graphs containing no circuit through some three prescribed elements (vertices or edges), the lack of such a circuit in $G - P$ is certified by what we refer to as an $\{x_3, x_4, x_5\}$ -frame (see DEFINITION A below).

§3.1 Frames. A pair (X, F) with $X \subset V(G)$ and $F \subseteq E(G - X)$ is called a *separator* if the number of components of $G - X - F$ is greater than the *capacity*

$$\gamma(X, F) := |X| + \sum \left\{ \lfloor \frac{1}{2} |\text{bnd}_{G-X} Q| \rfloor : Q \text{ is a component of } G[F] \right\}. \quad (3.3)$$

Here, $G[F] = (V(F), F)$, where $V(F)$ is the set of vertices incident with F . Given a set $A \subseteq V(G) \cup E(G)$, we specify (X, F) as an *A-separator* if A meets $> \gamma(X, F)$ components of $G - X - F$. A pair (G, A) cannot admit both an *A-circuit* and an *A-separator*. In general, neither of these exists, except for quite special cases when the “circuit-separator alternative” holds. The simplest such case is the generalization of the well-known theorem of Whitney (case $A \subseteq V(G)$, $|A| = 2$) proved by Mesner and Watkins [5].

Throughout the remainder of this section $\kappa(G) = 2$ and $U \subset E(G) \cup V(G)$, $|U| = 3$. The 2-hammocks of G satisfying $|U \cap H| \leq 1$ are referred to as *U-hammocks*. For two collections $\mathbf{L}, \mathbf{L}'$ of *U-hammocks*, let $\mathbf{L} \preceq \mathbf{L}'$ mean that each $H \in \mathbf{L}$ is a subgraph of some $H' \in \mathbf{L}'$, and let $\mathbf{L}$ be called *maximal* if $\mathbf{L} \preceq \mathbf{L}'$ implies $\mathbf{L} = \mathbf{L}'$. In a maximal collection, a common vertex of two hammocks is an end of each.

DEFINITION A.

A pair $\mathcal{X} = (X, F)$, where $X \subset V(G)$ and $F \subset E(G - X)$, is called a *U-frame* if the subgraph $G(\mathcal{X}) := (X \cup V(F), F)$ has exactly two components, and G is the union of $G(\mathcal{X})$ and a collection $\mathcal{B}$ of *U-hammocks* such that:

(A.1) a member of $\mathcal{B}$ has its ends in distinct components of $G(\mathcal{X})$;

(A.2) $U \cap H \neq \emptyset$ holds for three $H \in \mathcal{B}$; and $u \in \text{int}H$ of some $H \in \mathcal{B}$ for each vertex $u \in U$.

(A.3) $0 \leq |X| \leq 2$; and if $X = \emptyset$ then $|\mathcal{B}| = 3$.

(A.4) $\mathcal{B}$ is maximal subject to (A.1-3).

The members of $\mathcal{B}$ will be referred to as $\mathcal{X}$ -*hammocks*. Due to the maximality of $\mathcal{B}$, in case $F \neq \emptyset$:

(A.5) the ends of any two $\mathcal{X}$ -hammocks in the same component of $G[F]$ are distinct;

(A.6) the components of $G[F]$ are 2-connected and of order ≥ 3 .

We say that a U -frame $\mathcal{X}$ *separates* U . Figure 1 illustrates the 3 typical forms of a frame. The $\mathcal{X}$ -hammocks meeting U are called *seeded* and the members of U are called *seeds*. A




Figure 1: Frames $\mathcal{X} = (X, F)$ separating $\{o, o, o\}$.

through path containing the seed of a seeded $\mathcal{X}$ -hammock is called a *seeded-through path*; the latter exists due to (2.1). If x is a vertex in a seeded $\mathcal{X}$ -hammock, then a seeded-through path containing x is called a *seeded-through- x -path* (such paths do not necessarily exist).

Since any nontrivial $\mathcal{X}$ -hammock H becomes 2-connected once an ear connecting its ends is appended, H , if not 2-connected, is a chain of blocks. Consequently, if H is seeded with seed x , has ends z, z' , and $y \in \text{int}H - x$, then H contains a $y\ell$ -path P satisfying $x \in \text{int}P$, $\ell \in \{z, z'\}$. Such a path is called a *seeded- y -path*. The characterization of the 2-connected graphs containing no circuit traversing 3 prescribed elements of the graph (edges or vertices) is given by the following.

3.4. G has no U -circuit if and only if G has a U -frame.

Theorem 3.4 is derivable from the (elegant) fact that: a 3-connected (not necessarily simple) graph (V, E) has no A -circuit, where $A \subset V \cup E$ and $|A| = 3$, if and only if A consists of edges, and A is either an edge-cut, a claw ($K_{1,3}$), or contains two parallel edges.

Ma and Yu [2] use a close variant of the notion of frames; their variant, they explain, can be derived from the Mesner-Watkins argument [5]. The difference between Ma and Yu's variant to ours is that they ignore unseeded U -hammocks and essentially work with pairs $\mathcal{X} = (X, F)$ where $G(X)$ may have a single component instead of precisely 2; also in their version, U is always a set of vertices.

§3.2 Proof of 3.2. The reader should be reminded of the AGREEMENT specified in §2. Throughout this section, G is 5-connected and contains no TK_5 , P is as in 3.2, $U = \{x_3, x_4, x_5\}$ as these are defined in (3.1), $v \in \text{int}P$ and for such a v , U_v denotes a set of size 3 containing v satisfying $|U \cap U_v| = 2$. Also, if $\mathcal{X}$ is a Y -frame in graph J separating some 3-element set $Y \subset V(J) \cup E(J)$, then the components of $J(\mathcal{X})$ are referred to as F_1 and F_2 , and the end of an $\mathcal{X}$ -hammock H contained in F_i is denoted $z_{i,H}$, for $i = 1, 2$.

To reach a contradiction, we show that a U_v -circuit satisfying (F.2) exists. To this end, suffices to show that there is an

$$\ell\ell'\text{-path } Q \text{ in } G - P \text{ satisfying } \ell, \ell' \in N_{G-P}(v) \text{ and } |V(Q) \cap U| = 2. \quad (3.5)$$

Consequently, throughout this section, we assume, towards contradiction, that a path satisfying (3.5) does not exist.

By (F.1) and 3.4, there is a U -frame in $G - P$. In what follows, we study U -frames in $G - P$ (under the assumption that a path satisfying (3.5) does not exist).

REDUCTION STEPS. Let J be a graph with $\kappa(J) = 2$ and let $\mathcal{X}$ be a W -frame for some $W \subset V(J)$, $|W| = 3$. Let H be a seeded $\mathcal{X}$ -hammock with seed x . If $y \in V(H)$, then an $(X_y = \{x, y, z_{1,H}, z_{2,H}\})$ -circuit in $R_H^1 = H + z_{1,H}z_{2,H}$, implies the existence of a seeded-through- y -path in H . Next, if $y, y' \in V(H)$, then an $(X_{y,y'} = \{x, yy', z_{1,H}, z_{2,H}\})$ -circuit in $R_H^2 = R_H^2(y, y') = H + z_{1,H}z_{2,H} + yy'$, implies the existence of a yy' -path in $G - P$ meeting precisely two seeds. Consequently, by 3.4:

(R.1) *If $y \in V(H)$ and R_H^1 contains no seeded-through- y -path, then there is an X_y -frame in R_H^1 ; and thus $y \in (\text{int}_{G-P}H) \setminus \{x\}$.*

(R.2) *If $y, y' \in V(H)$ and $G - P$ does not contain a yy' -path meeting precisely two seeds, then there is an $X_{y,y'}$ -frame in $R_H^2(y, y')$.*

For $i = 1, 2$, we refer to the frames defined in (R.i) as $\mathcal{X}_H = (X_H, F_H)$. The seeded $\mathcal{X}_H$ -hammocks containing $x, y, z_{1,H}, z_{2,H}$ in (R.1), and $x, yy', z_{1,H}, z_{2,H}$ in (R.2), are denoted by K_H, K'_H, K''_H , respectively. By $\mathcal{P}_H$ we mean a 2-linkage in $G - P$ connecting $\{z_{1,H}, z_{2,H}\}$ to the ends of K''_H ; such a linkage is clearly contained in $K''_H - z_{1,H}z_{2,H}$. Let $\ell_{1,H}, \ell_{2,H}$ be the ends of K''_H linked to $z_{1,H}, z_{2,H}$ in $\mathcal{P}_H$, respectively. Call the components of $R_H^i(\mathcal{X}_H)$, $i = 1, 2$, containing $\ell_{1,H}, \ell_{2,H}$, by $F_{1,H}$ and $F_{2,H}$, respectively.

CONSTRUCTIONS. We list several settings and constructions of paths therein which yield a path satisfying (3.5) and that rise frequently in subsequent arguments. Let J be a graph with $\kappa(J) = 2$ and let $\mathcal{X}$ be a W -frame for some $W \subset V(J)$, $|W| = 3$.

(S.1) *Let H, H' be seeded $\mathcal{X}$ -hammocks and let $y \in \text{int}_J H$. Let $\mathcal{X}_{H'}$ be an X_u -frame or an $X_{u,u'}$ -frame in $R_{H'}^i$ per (R.i) for $i = 1, 2$, where $u, u' \in V(H')$. If H contains a seeded-through- y -path, then J contains a $(y, \{u, u'\})$ -path meeting precisely two seeds.*

Proof. Let S be a seeded-through- y -path in H such that $z_{i,H}, y, x, z_{3-i,H}$ appear in this order along S for some $i = 1, 2$, and where x is the seed of H . A path satisfying the assertion is

contained in the union of the following paths: the $yz_{3-i,H}$ -subpath of S , a path in F_{3-i} linking the ends of H and H' , the $\ell_{3-i,H'}z_{3-i,H'}$ -path in $\mathcal{P}_{H'}$, a path in $F_{3-i,H'}$ linking the ends of $K_{H'}$ and $K_{H'}''$, a seeded-through path in $K_{H'}$, a path in $F_{i,H'}$ linking the ends of $K_{H'}$ and $K_{H'}'$, a path in $K_{H'}' - uu'$ linking $z_{i,K_{H'}'}$ to a member of $\{u, u'\}$ not meeting the other end of $K_{H'}'$. Easy to see that the last path exists. ■

(S.2) Let H, H' be seeded $\mathcal{X}$ -hammocks and let $y \in V(H), y' \in V(H')$. Let $\mathcal{X}_H$ and $\mathcal{X}_{H'}$ be an X_y -frame and an $X_{y'}$ -frame in R_H^1 and $R_{H'}^1$, respectively. Pick an $i \in \{1, 2\}$ and consider the following union of paths: a yz_{i,K_H} -path in K_H' not meeting z_{3-i,K_H} (such a path exists by (2.1) and $y \in \text{int}_J H$ by (R.1)), a path in $F_{i,H}$ linking the ends of K_H' and K_H'' , the $\ell_{i,H}z_{i,H}$ -path in $\mathcal{P}_H$, a path in F_i linking the ends of H and H'' , where H'' is the third seeded $\mathcal{X}$ -hammock, a seeded-through path in H'' , a path in F_{3-i} linking the ends of H' and H'' , the $z_{3-i,H'}\ell_{3-i,H'}$ -path in $\mathcal{P}_{H'}$, a path in $F_{3-i,H'}$ connecting the ends of $K_{H'}$ and $K_{H'}''$, a seeded-through path in $K_{H'}$, a path in $F_{i,H'}$ linking the ends of $K_{H'}$ and $K_{H'}'$, a $z_{i,K_{H'}'}y'$ -path in $K_{H'}'$ not meeting $z_{i,K_{H'}'}$.

The resulting subgraph R is called a *trail*. Such is a yy' -path in J meeting precisely two seeds provided that $R \cap F_i$ and $R \cap F_{i,H'}$ are disjoint. By the definition of frames, this does not occur provided F_i and $F_{i,H'}$ coincide into a singleton.

Due to symmetry, the above construction defines 4 distinct trails: indeed we may choose to start the construction from either y or y' (here we started from y) for each such choice we may choose $i \in \{1, 2\}$.

(S.3) Let $y \in \text{int}_J H, y' \in \text{int}_J H'$, where H, H' are a seeded and an unseeded $\mathcal{X}$ -hammocks, respectively. If H contains a seeded-through- y -path, then J contains a yy' -path meeting precisely two seeds.

Proof. Let S be a seeded-through- y -path in H such that $z_{i,H}, y, x, z_{3-i,H}$ appear in this order along S for some $i = 1, 2$, and where x is the seed of H . A path satisfying the assertion is contained in the union of the following paths: the $yz_{3-i,H}$ -subpath of S , a path in F_{3-i} linking the ends of H and H'' , where H'' is some seeded $\mathcal{X}$ -hammock, a seeded-through path in H'' , a path in F_i linking the ends of H'' and H' , a $z_{i,H'}y'$ -path in H' not meeting $z_{3-i,H'}$. ■

(S.4) Let $y \in V(H), y' \in \text{int}_J H'$, where H and H' are a seeded and unseeded $\mathcal{X}$ -hammocks, respectively. Let $\mathcal{X}_H$ be an X_y -frame in R_H^1 . Pick an $i \in \{1, 2\}$ and consider the following union of paths: a yz_{i,K_H} -path in K_H' not meeting $z_{3-i,K_H'}$, a path in $F_{i,H}$ linking the ends of K_H and K_H' , a seeded-through path in K_H , a path in $F_{3-i,H}$ linking the ends of K_H and K_H'' , the $\ell_{3-i,H}z_{3-i,H}$ -path in $\mathcal{P}_H$, a path in F_{3-i} linking the ends of H and H'' where H'' is some seeded $\mathcal{X}$ -hammock, a seeded-through path in H'' , a path in F_i linking the ends of H'' and H' , a $z_{i,H'}y'$ -path in H' not meeting $z_{3-i,H'}$.

The resulting subgraph R is called a *poor trail*. Such is a yy' -path in J meeting precisely two seeds provided that $R \cap F_i$ and $R \cap F_{i,H}$ are disjoint. By the definition of frames, this does not occur provided F_i and $F_{i,H}$ coincide into a singleton.

Due to symmetry, the above construction defines 2 distinct poor trails as we may choose $i \in \{1, 2\}$.

PROPERTIES OF U -FRAMES IN $G - P$. Let $\mathcal{X}$ be a U -frame in $G - P$, $\mathcal{B}$ its collection of $\mathcal{X}$ -hammocks, and let $v \in \text{int}P$. We consider several properties of $\mathcal{X}$ and $N_{G-P}(v)$.

$$(3.2.A) \quad N_{G-P}(v) \subseteq \bigcup_{H \in \mathcal{B}} V(H).$$

Proof. The following two claims (3.2.A.1) and (3.2.A.2) are sufficient.

(3.2.A.1) *If $|N_G(v) \cap V(F_i)| \geq 2$, for some $i = 1, 2$, then a U_v -circuit satisfying (F.2) exists.*

Proof. Let $y, y' \in N_G(v) \cap V(F_i)$, for some $i = 1, 2$. Let $\mathcal{P}$ be a 2-linkage in F_i connecting $\{y, y'\}$ and two ends in F_i of some two seeded $\mathcal{X}$ -hammocks, namely H and H' . Let Y and Y' be seeded-through paths in H and H' , respectively. Let Q be a path in F_{3-i} connecting the ends of H and H' in F_{3-i} . Clearly, $\mathcal{P} \cup Y \cup Y' \cup Q \cup \{yv, y'v\}$ is a U_v -circuit satisfying (F.2). $\square$

By (3.2.A.1), and since $|N_{G-P}(v)| \geq 3$ (as P is induced), and since $G(\mathcal{X})$ has precisely two components, $N_H(v) \neq \emptyset$ for at least one $H \in \mathcal{B}$. To conclude then, we prove the following.

(3.2.A.2) *If $N_G(v) \cap \text{int}_{G-P}F_i \neq \emptyset$, for some $i = 1, 2$, and $N_H(v) \neq \emptyset$, for some $H \in \mathcal{B}$, then a U_v -circuit satisfying (F.2) exists.*

Proof. Here we prefer to use the following notation. Let H_i denote the seeded $\mathcal{X}$ -hammocks and let z_i^j denote the end of H_i in F_j , for $i = 1, 2, 3$ and $j = 1, 2$. Let S_i denote a seeded-through path in H_i , and let $Q_{i,x}$ denote a seeded- x -path in H_i , where $x \in N_{H_i}(v)$. By $P_{k,\ell}^i$ we mean a $z_k^i z_\ell^i$ -path in F_i , for $i = 1, 2$ and $k, \ell = 1, 2, 3$, $k \neq \ell$. Let $y \in N_G(v) \cap \text{int}_{G-P}F_r$, for some $r = 1, 2$. By $F_{k,\ell}$ we mean a $(y, \{z_k^r, z_\ell^r\})$ -2-fan in F_r , for $r = 1, 2$, $k, \ell = 1, 2, 3$, $k \neq \ell$. Let $\mathcal{P}_{j,k,\ell}$ denote a $(\{y, z_j^r\}, \{z_k^r, z_\ell^r\})$ -2-linkage in F_r . We observe the following.

(a) If $x \in N_{H_j}(v)$, $j = 1, 2, 3$, then $Q_{j,x}$ ends in z_j^r . Otherwise, a path satisfying (3.5) is contained in $Q_{j,x} \cup F_{j,k} \cup S_k \cup P_{j,k}^{3-r}$, where $j \neq k$, $j, k = 1, 2, 3$.

(b) By (a) and (3.2.A.1), if $N_{H_j}(v) \neq \emptyset$, for some $j = 1, 2, 3$, then $N_{F_{3-r}}(v) \cap \text{int}_{G-P}F_{3-r} = \emptyset$.

(c) If $x \in N_{H_j}(v)$, then $N_{H_k}(v) = N_{H_\ell}(v) = \emptyset$, where $\{j, k, \ell\} = \{1, 2, 3\}$. To see this, let $z \in N_{H_k}(v)$. By (a), $Q_{j,x}$ and $Q_{k,z}$ end at z_j^r and z_k^r , respectively. Thus, a path satisfying (3.5) is contained in $Q_{j,x} \cup Q_{j,z} \cup P_{j,k}^r$.

(d) If $x, z \in N_{H_j}(v)$, $j = 1, 2, 3$, then a path satisfying (3.5) exists. Let $\mathcal{P} \subseteq H_j$ be an $(\{x, z\}, \{z_j^r, z_j^{3-r}\})$ -2-linkage. The seed of H_j is not contained in $\mathcal{P}$. Indeed, if so, a path satisfying (3.5) is contained in $\mathcal{P} \cup P_{j,k}^r \cup P_{j,k}^{3-r} \cup S_k$, where $j \neq k = 1, 2, 3$. Consequently, a path satisfying (3.5) is contained in $R \cup \mathcal{P}_{j,k,\ell} \cup S_k \cup P_{k,\ell}^{3-r} \cup S_\ell$, where $R \in \mathcal{P}$ and has z_j^r as an end and $\{j, k, \ell\} = \{1, 2, 3\}$.

(e) If $x \in N_{G-P}(v) \cap \text{int}_{G-P}H$, where $H \in \mathcal{B}$ and unseeded, then a path satisfying (3.5) exists. To see this, let $R \subset H$ be an xw -path such that w is the end of H in F_r and such that R meets only one end of H ; by (2.1), R exists. Let $\mathcal{P}$ be a $(\{w, y\}, \{z_k^r, z_\ell^r\})$ -2-linkage in F_r . A path satisfying (3.5) is contained in $R \cup \mathcal{P} \cup S_\ell \cup S_k \cup P_{k,\ell}^{3-r}$, where $k, \ell = 1, 2, 3$, $k \neq \ell$.

To conclude the proof consider the following. We have that $N_H(v) \neq \emptyset$ for some $H \in \mathcal{B}$, by (3.2.A.1). Presence of an unseeded $\mathcal{X}$ -hammock implies that at least one of the compo-

nents of $G'(\mathcal{X})$, where $G' = G - P$, is a singleton. In our setting, F_{3-r} will be a singleton. Thus, by (3.2.A.1) and (b), (e), we may assume that H is seeded, regardless of the existence of unseeded $\mathcal{X}$ -hammocks. Now, as $N_{F_{3-r}}(v) \cap \text{int}_{G-P} F_{3-r} = \emptyset$ and $|N_{F_r}(v)| = 1$, by (b) and (3.2.A.1), respectively, then either $|N_H(v)| \geq 2$ contradicting (d), or $N_{G-P}(v)$ meets at least two seeded hammocks contradicting (c), or $N_{G-P}(v)$ meets the interior of an unseeded $\mathcal{X}$ -hammock contradicting (e). $\square$

(3.2.B) *If $y \in N_{G-P}(v) \cap \text{int}_{G-P} H$ where H is a seeded $\mathcal{X}$ -hammock containing a seeded-through- y -path, then no other seeded $\mathcal{X}$ -hammock meets ≥ 2 members of $N_{G-P}(v)$.*

Proof. Let $u, u' \in N_{H'}(v)$, where $H' \neq H$ is a seeded $\mathcal{X}$ -hammock. Clearly $u, u' \neq y$. By (R.2), there is an $X_{u,u'}$ -frame in $R_{H'}^2(u, u')$. Thus a path satisfying (3.5) exists, by (S.1). $\blacksquare$

(3.2.C) *Let $y \in N_H(v)$, $y' \in N_{G-P}(v) \cap \text{int}_{G-P} H'$, where H, H' are seeded $\mathcal{X}$ -hammocks. If there is no seeded-through- y -path in H and H' has a seeded-through- y' -path, then a path satisfying (3.5) exists.*

Proof. By (R.1), there is an X_y -frame in R_H^1 . Thus a path satisfying (3.5) exists, by (S.1). $\blacksquare$

PROPERTIES OF MINIMUM U -FRAMES. For a U -frame $\mathcal{X}$ in $G - P$, put

$$\alpha(\mathcal{X}) = |\bigcup \{V(H) : H \text{ is a seeded } \mathcal{X}\text{-hammock}\}|. \quad (3.6)$$

We write $\mathcal{X}_1 \preceq \mathcal{X}_2$ if $\alpha(\mathcal{X}_1) \leq \alpha(\mathcal{X}_2)$, where $\mathcal{X}_1, \mathcal{X}_2$ are U -frames in $G - P$. We say that a U -frame $\mathcal{X}$ is *minimum* if $\mathcal{X}' \preceq \mathcal{X}$ implies $\mathcal{X}' = \mathcal{X}$ for every U -frame $\mathcal{X}'$.

(3.2.D) *Let $\mathcal{X}$ be a minimum U -frame. If $y \in N_H(v)$, $y' \in N_{H'}(v)$, where H, H' are seeded $\mathcal{X}$ -hammocks, then H, H' contain a seeded-through- y -path and a seeded-through- y' -path, respectively.*

Proof. At least one of y, y' is contained internally in H, H' , respectively; otherwise the claim follows trivially from (2.1). Consequently, by (3.2.C), we may assume that H, H' contain no seeded-through- y -path and no seeded-through- y' -path, respectively. Thus, by (R.1), there are an X_y -frame and an $X_{y'}$ -frame in R_H^1 and $R_{H'}^1$, respectively.

The assumption that there is no path satisfying (3.5) implies that all (symmetric) trails produced by (S.2) fail to provide such a path. This occurs provided $F_i, F_{i,H}, F_{i,H'}$ all coincide into a singleton q_i , for $i = 1, 2$. The pair $\mathcal{X}' = (\{q_1, q_2\}, \emptyset)$ then is a U -frame with a collection $\mathcal{B}'$ of U -hammocks satisfying $K_H, K_{H'}, H'' \in \mathcal{B}'$ implying that $\alpha(\mathcal{X}') < \alpha(\mathcal{X})$ since $K_H \subset H, K_{H'} \subset H'$. $\blacksquare$

(3.2.E) *Let $\mathcal{X}$ be a minimum U -frame. If $N_H(v), N_{H'}(v) \neq \emptyset$ and $N_H(v) \neq N_{H'}(v)$ for some two seeded $\mathcal{X}$ -hammocks $H, H' \in \mathcal{B}$, then $N_{G-P}(v) \subseteq V(H) \cup V(H')$.*

Proof. Let $y \in N_H(v), y' \in N_{H'}(v), y \neq y'$. Let x, x' be the seeds of H, H' , respectively. Seeded-through- b -paths, $b \in \{y, y'\}$ in H and H' , respectively, exist by (3.2.D). Such have

noncorresponding orderings in the sense that: if H has a seeded-through- y -path such that $z_{i,H}, y, x, z_{3-i,H}$ appear in this order along the path, then a seeded-through- y' -path in H' has $z_{i,H'}, x', y', z_{3-i,H'}$ appear in this order along it. Indeed, if not so, and the orderings correspond, then since $y \neq y'$, a path satisfying (3.5) clearly exists. Consequently, $N_{G-P}(v)$ does not meet the interior of the third seeded $\mathcal{X}$ -hammock.

Let then $H'' \in \mathcal{B}$ be unseeded such that $y'' \in N_{G-P}(v) \cap \text{int}_{G-P}H''$. By the definition of frames, F_i is a singleton for some $i = 1, 2$. Consequently, at least one of $\{y, y'\}$, say y , is contained internally in its corresponding $\mathcal{X}$ -hammock. A path satisfying (3.5) then exists by (S.3). Such an H'' exists by (3.2.A) and F_i being a singleton for some $i = 1, 2$. ■

(3.2.F) *Let $\mathcal{X}$ be a minimum U -frame. If $N_{G-P}(v) \cap \text{int}_{G-P}H \neq \emptyset$ for some seeded $\mathcal{X}$ -hammock H , then $N_{G-P}(v) \subseteq V(H)$.*

Proof. By (3.2.E), if $N_{G-P}(v) \cap \text{int}_{G-P}H' \neq \emptyset$ for some seeded $\mathcal{X}$ -hammock distinct of H , then $N_{G-P}(v) \subseteq V(H) \cup V(H')$. This implies that H, H' meet at least three distinct members of $N_{G-P}(v)$. A path satisfying (3.5) then exists, by (3.2.B) and (3.2.D).

Let H' be unseeded then. Thus, H has no seeded-through- y -path, $y \in N_{G-P}(v) \cap \text{int}_{G-P}H$; otherwise a path satisfying (3.5) exists by (S.3). Thus, by (R.1), there is an X_y -frame in R_H^1 . The assumption that no path satisfying (3.5) implies that all (symmetric) poor trails produced by (S.4) fail to provide such a path. This occurs provided F_i and $F_{i,H}$ coincide into a singleton q_i for $i = 1, 2$. Consequently, $\mathcal{X}' = (\{q_1, q_2\}, \emptyset)$ is a U -frame with a collection $\mathcal{B}'$ of U -hammocks satisfying $K_H, H'', H''' \in \mathcal{B}'$, where H'', H''' are the remaining seeded $\mathcal{X}$ -hammocks. As $K_H \subset H$, it follows that $\alpha(\mathcal{X}') < \alpha(\mathcal{X})$. ■

CONSEQUENCES. We learn the following from minimum U -frames in $G - P$.

(3.2.G) *A seed is not adjacent to $\text{int}P$.*

Proof. Let $x \in U$ be adjacent to $\text{int}P$ and let $\mathcal{X}$ be a minimum U -frame with H the $\mathcal{X}$ -hammock containing x . Let $v \in \text{int}P$ such that $x \in N_H(v)$. As $x \in \text{int}_{G-P}H$, by definition of frames, we have that $N_{G-P}(v) \subseteq V(H)$, by (3.2.F). Let $y \in N_H(v) \setminus \{x\}$. A $\{yx, z_{1,H}z_{2,H}\}$ -circuit clearly exists in $H + \{yx, z_{1,H}z_{2,H}\}$. Such a circuit is easily extended into a yx -path satisfying (3.5). ■

Property (3.2.G) implies that the seeded $\mathcal{X}$ -hammocks of a U -frame $\mathcal{X}$ have order ≥ 4 each. Thus, 5-connectivity then asserts that the interior of such hammocks must be adjacent to $\text{int}P$. Property (3.2.F) then implies that if $H_i, i = 1, 2, 3$, are the seeded $\mathcal{X}$ -hammocks of a minimum $\mathcal{X}$, then there is a partition $L_1 \cup L_2 \cup L_3 \cup L_4 = \text{int}P$ such that $L_i, i = 1, 2, 3$, consists of the vertices of $\text{int}P$ adjacent to $\text{int}_{G-P}H_i$, and L_4 is the complement of $L_1 \cup L_2 \cup L_3$. We have seen that for $i = 1, 2, 3$, $L_i \neq \emptyset$; moreover, if $v \in L_i$, then $N_{G-P}(v) \subseteq V(H_i)$, by (3.2.F).

Let $v \in L \in \{L_1, L_2, L_3\}$ and let H denote the seeded $\mathcal{X}$ -hammock corresponding to L . Clearly, (R.2) applies to H and any two members of $N_H(v)$; or a path satisfying (3.5) exists. Let $L_{v,H}$ denote the $\mathcal{X}_H$ -frames of type (R.2) in R_H^2 where (R.2) is applied to the seed of H and two members of $N_H(v)$; we have just seen that $L_{v,H} \neq \emptyset$. For $y \in V(H)$, that is not the seed of H , let $L_{y,H} = \emptyset$ if H contains a seeded-through- y -path. Otherwise let $L_{y,H}$ denote the $\mathcal{X}_H$ -frames of type (R.1) in R_H^1 , where (R.1) is applied to y and the seed of

H . Put $\mathbf{L}_H = (\bigcup_{v \in L} \mathbf{L}_{v,H}) \cup (\bigcup_{y \in V(H)} \mathbf{L}_{y,H})$. For $\mathcal{X}_H \in \mathbf{L}_H$, put $\beta(\mathcal{X}_H) = |V(K_H)|$ and let $\beta = \min\{\beta(\mathcal{X}_H) : \mathcal{X}_H \in \mathbf{L}_H\}$.

We are now ready to prove **3.2** and consequently **1.3**.

Proof of 3.2. Let $\mathcal{X}$ be a minimum U -frame in $G - P$ and let L_i , $1 \leq i \leq 4$ be the partition of $\text{int}P$ induced by $\mathcal{X}$ as defined above. Let $L \in \{L_1, L_2, L_3\}$, let H be the seeded $\mathcal{X}$ -hammock corresponding to L , and let x be its seed.

$$\text{Choose } \mathcal{X}_H \in \mathbf{L}_H \text{ satisfying } \beta(\mathcal{X}_H) = \beta. \quad (3.7)$$

The minimality of $\mathcal{X}_H$ as in (3.7) implies that for every $v' \in L$ and $y \in V(K_H) \setminus \{x\}$,

$$(i) |N_{K_H}(v')| \leq 1 \text{ and } (ii) K_H \text{ contains a seeded-through-}y\text{-path.} \quad (3.8)$$

Proof. To see (3.8), let ℓ, ℓ' denote the ends of K_H . Observe that an $\{x, yy', \ell \ell'\}$ -circuit in $K_H + \{yy', \ell \ell'\}$, for some two $y, y' \in N_{K_H}(v')$, easily extends into a yy' -path satisfying (3.5). Consequently, if (3.8) (i) is not satisfied, then an $\{x, yy', \ell \ell'\}$ -frame $\mathcal{X}_{K_H}$ exists in $K_H + \{yy', \ell \ell'\}$, by (R.2). In a similar manner, if K_H contains no seeded-through- y -path, then $K_H + \ell \ell'$ has an $\{x, y, \ell \ell'\}$ -frame $\mathcal{X}_{K_H}$ by in (R.1).

Assume now, to the contrary, that (3.8) (i) or (ii) is not satisfied for some $v' \in L$ and $u, u' \in N_{K_H}(v')$ or some $y \in V(K_H) \setminus \{x\}$, respectively, and let $\mathcal{X}_{K_H}$ be a frame in $R_{K_H}^2(u, u')$ and $R_{K_H}^1$, respectively. Consider the following frame $\mathcal{Y}$. Let $G' \in \{R_{K_H}^1, R_{K_H}^1(u, u')\}$ and put: $G'(\mathcal{Y}) = G'(\mathcal{X}_{K_H}) = F_{1,K_H} \cup F_{2,K_H}$, and the $\mathcal{Y}$ -hammocks are K_{K_H}, K'_{K_H} , and the third $\mathcal{Y}$ -hammock is the union of $K''_{K_H}, F_{1,H}, F_{2,H}, K'_H$, and K''_H .

If $\mathcal{X}_{K_H}$ is the result of a violation of (3.8) (i) for some $v' \in L$ and $u, u' \in N_{K_H}(v')$, then $\mathcal{Y} \in \mathbf{L}_{v',H}$ (and thus in $\mathbf{L}_H$). If $\mathcal{X}_{K_H}$ is the result of a violation of (3.8) (ii) for some $y \in V(K_H) \setminus \{x\}$, then $\mathcal{Y} \in \mathbf{L}_{y,H}$ (and thus in $\mathbf{L}_H$).

Property (3.8) now follows since $K_{K_H} \subset K_H$ implying that $\beta(\mathcal{Y}) < \beta(\mathcal{X}_H)$; contradicting (3.7). $\square$

By (3.2.G), $|V(K_H)| \geq 4$; thus, 5-connectivity and (3.8) then imply that there are a v' and a y satisfying $v' \in L$ and a $\{y\} = \text{int}_{G-P} K_H \cap N_H(v')$. As $N_{G-P}(v') = N_H(v')$, then $N_H(v') \setminus \{y\} \subseteq H - \text{int}_{G-P} K_H$, by (3.2.F), (3.2.G) and (3.8).

To conclude the proof, suffices that we show that

$$\text{there are } i = 1, 2 \text{ and } y', y'' \text{ such that } y', y'' \in V(F_{i,H}) \cap (N_H(v') \setminus \{y\}). \quad (3.9)$$

Indeed, if so, it is easy to see that: a $(\{y', y''\}, \{z_{i,K_H}, \ell_{i,H}\})$ -2-linkage in $F_{i,H}$, the linkage $\mathcal{P}_H$, a seeded-through path in K_H , and a seeded-through path in H' , where H' is some seeded $\mathcal{X}$ -hammock, can be extended into a $y'y''$ -path satisfying (3.5).

To establish (3.9), we argue as follows. Let $y' \in N_H(v') \setminus \{y\}$. Let $S \subseteq K_H$ be the seeded-through- y -path in K_H , which exists by (3.8), such that $z_{i,K_H}, y, x, z_{3-i,K_H}$ appear in this order along S , where x is the seed of K_H . Suffices that we argue the following:

(a) $y' \notin F_{i,H} \cup (K'_H - z_{3-i,K'_H})$. Indeed, if this is not so, then a path satisfying (3.5) is contained in the union of the following paths: the yz_{3-i,K_H} -subpath of S , a path in $F_{3-i,H}$ linking the

ends of K_H and K_H'' , the members of $\mathcal{P}_H$, a path in F_{3-i} linking the ends of H and H' , where H' is some seeded $\mathcal{X}$ -hammock, a seeded-through path in H' , a path in F_i linking the ends of H' and H , a path in $F_{i,H} \cup (K_H' - \{z_{3-i,K_H'}\})$ linking y' with $\ell_{i,H}$.

(b) $y' \notin \text{int}_{G-P} K_H''$. If not so, then a $(\{y', \ell_{3-i,H}\}, \{z_{1,H}, z_{2,H}\})$ -2-linkage is clearly contained in $K_H'' - z_{1,H} z_{2,H}$ and is easily extended into a path satisfying (3.5). ■

§4 Apex graphs containing $K_{2,3}$. To outline our approach for proving **1.2**, suppose now that G is a 5-connected nonplanar apex graph containing no TK_5 and $K \subseteq G$, where K is as in (3.1). By **1.1**, we may assume throughout this section that

$$\{x_1, x_2\} \text{ and } \{x_3, x_4, x_5\} \text{ are stable.} \quad (4.1)$$

By **4.9**, there is an induced $x_1 x_2$ -path P such that $x_3, x_4, x_5 \notin V(P)$ and with a P -bridge meeting at least two 2-valent vertices of K . We propose that two cases be examined:

CONNECTED CASE: P can be chosen so that there is a P -bridge containing $\{x_3, x_4, x_5\}$, or
DISCONNECTED CASE: such a choice is impossible.

In the CONNECTED CASE, by **4.7** with $Y = \{x_3, x_4, x_5\}$ and $X = \emptyset$ (X, Y - see formulation of **4.7**), P can be chosen such that

$$G - P \text{ is connected with each leaf block meeting } \{x_3, x_4, x_5\} \text{ internally.} \quad (4.2)$$

By **1.3** we may assume that $\kappa(G - P) = 1$ and thus $G - P$ is a chain or a claw-chain of blocks. By **4.14**, assuming that $TK_5 \not\subseteq G$ implies that neither x_1 nor x_2 is an apex vertex of G . To reach contradiction in this case we that

4.3. *if $\kappa(G - P) = 1$, each leaf block of $G - P$ meets $\{x_3, x_4, x_5\}$ internally, and $a \in V(G) \setminus \{x_1, x_2\}$ is an apex vertex of G , then $TK_5 \subseteq G$.*

For the DISCONNECTED CASE we argue the following. By **4.9**, **1.3** and **4.3**, we may assume, without loss of generality, that P can be chosen so that there is P -bridge containing $\{x_4, x_5\}$, and there is no such P with a P -bridge containing $\{x_3, x_4, x_5\}$. Thus, by **4.7** with $Y = \{x_4, x_5\}$ and $X = \{x_3\}$, P can be chosen so that $G - P$ consists of precisely two components:

$$x_3 \text{ and a chain of blocks } H \text{ with each leaf block meeting } \{x_4, x_5\} \text{ internally.} \quad (4.4)$$

If $\kappa(H) \geq 2$, we utilize the assumption that (F.2) fails and use **3.4** to prove that

4.5. *if $\kappa(H) \geq 2$ and $a \in V(G) \setminus \{x_1, x_2\}$ is an apex vertex of G , then $TK_5 \subseteq G$.*

To conclude our arguments for the DISCONNECTED CASE, and indeed our proof of **1.2**, we show that

4.6. *if $\kappa(H) = 1$ and $a \in V(G) \setminus \{x_1, x_2\}$ is an apex vertex of G , then $TK_5 \subseteq G$.*

In the remainder of this section we prove **4.3**, **4.5**, and **4.6**.

§4.1 Rooted nonseparating paths. The purpose of this section is to prove **4.7** and **4.9**; usage of these is outlined above. Lemma **4.7** is modeled after [2, Lemma 2.3] and serves as a generalized variant of [2, Lemma 2.3]. Our proof of **4.7** follows from the three claims: **(4.7.A)**, **(4.7.B)**, and **(4.7.C)** listed below. The proof of **(4.7.A)** is precisely the proof of an analogous claim that Ma and Yu use to prove [2, Lemma 2.3]; we do not see a need to change their argument. To prove **(4.7.B)**, we use the approach of Ma and Yu for an analogous claim in their argument, but the implementation of their approach here requires adjustments resulting from the differences between **4.7** and [2, Lemma 2.3]. Claim **(4.7.C)** is new. Consequently, we attribute the following to Ma and Yu, but see no option but to include a complete proof of **4.7**.

4.7. Suppose $\kappa(G) \geq 5$. Let $X, Y \subseteq V(G)$ be disjoint such that Y is stable and $|Y| \geq 2$ (possibly, $X = \emptyset$). Let $x, y \in V(G) \setminus (X \cup Y)$. Suppose the following terms are satisfied:

- (4.7.a)** there is an induced xy -path in $G - (X \cup Y)$ with a bridge containing Y , and
- (4.7.b)** there is no induced xy -path in $G - (X \cup Y)$ with a bridge containing $Y \cup \{x\}$ for any $x \in X$.

Then, there is an induced xy -path P in $G - (X \cup Y)$ such that:

- (4.7.c)** each P -bridge either meets X or contains Y , and
- (4.7.d)** the interior of each leaf block of the P -bridge containing Y meets Y , and
- (4.7.e)** if $|X| < \kappa(G) - 2$, then each P -bridge meeting X is a singleton.

Proof. By **(4.7.a)** and **(4.7.b)**, there exist 4-tuples $(P, B, H, \mathcal{B})$ where P is an induced xy -path with a P -bridge B containing Y , $H \subseteq B$ contains Y with each of its leaf blocks meeting Y internally, and $\mathcal{B}$ is the union of P -bridges meeting X . Existence of H follows from the fact that given a subgraph of B containing Y one may trim the block tree of such a subgraph by successively removing the interior of leaf blocks not meeting Y internally.

Choose a 4-tuple $\mathcal{X} = (P, B, H, \mathcal{B})$ such that

- (I) H is maximal,
- (II) subject to (I) $\mathcal{B}$ is maximal, and
- (III) subject to (I) and (II) the number of P -bridges not meeting $X \cup Y$ is minimum.

Claims **(4.7.A,B,C)**(see below) imply **4.7**.

(4.7.A) In $\mathcal{X}$: there are no P -bridges not meeting $X \cup Y$.

Proof. Assume, towards contradiction, that there is a P -bridge D not meeting $X \cup Y$ and let v_1, v_2 be its extremal attachments on P . Let $C \neq D$ be a P -bridge adjacent to $(v_1 P v_2)$; existence of which follows from 5-connectivity.

For an induced $v_1 v_2$ -path P' in $G[V(D) \cup \{v_1, v_2\}]$, let $\mathcal{D}$ denote the union of the P' -bridges of D not adjacent to $(v_1 P v_2)$. Let $\mathcal{D}'$ denote the remaining P' -bridges of D . Choose P' so that $\mathcal{D}$ is minimal and let P'' be the induced xy -path obtained from P by replacing $(v_1 P v_2)$ with P' .

$\mathcal{D}$ is nonempty; for otherwise $(D - P'') \cup C \cup (v_1 P v_2)$ is contained in some P'' -bridge contradicting (III). Let then $D_1, \dots, D_k$ be the components of $\mathcal{D}$, and let a_i, b_i be the extremal attachments of D_i on P'' , for $1 \leq i \leq k$. Let $c_i \in (a_i P'' b_i)$ adjacent to $D - (P'' \cup D_i)$ or $(v_1 P v_2)$; existence of c_i follows from 5-connectivity for otherwise $\{a_i, b_i, v_1, v_2\}$ is a 4-disconnector of G .

The vertex c_i is not adjacent to a component of $\mathcal{D}'$ or to $(v_1 P v_2)$ for then the induced

v_1v_2 -path in $G[V(D) \cup \{v_1, v_2\}]$ obtained from P' by replacing $a_iP'b_i$ with an induced a_ib_i -path in $G[V(D_i) \cup \{a_i, b_i\}]$ contradicts the choice of P' . Consequently, since P'' is induced, $N((a_iP''b_i)) \subseteq V(\mathcal{D})$, for $1 \leq i \leq k$. Therefore, $\bigcup_{i=1}^k (a_iP''b_i)$ is a subpath of P'' ; let a and b be its ends. Since $\{a, b, v_1, v_2\}$ is not a 4-disconnector of G , there exists a $c \in (aP''b)$ adjacent to a component of $\mathcal{D}'$ or to (v_1Pv_2) . Consequently, there is an $1 \leq i \leq k$ such that $c \in (a_iP''b_i)$ which is a contradiction. $\square$

(4.7.B) *In $\mathcal{X}$: H coincides with B .*

Proof. The H -bridges of $G - P$ are not adjacent to and do not meet members of $\mathcal{B}$, by (4.7.b). Consequently, the H -bridges of $G - P$ are subgraphs of B , by (4.7.A). There are no H -bridges of B with ≥ 2 attachments in H for the union of H and an H -ear in B linking two such attachments contradicts (I) as the property of each leaf block of such a union meeting Y is sustained.

It remains to prove that there are no H -bridges of B with a single attachment in H . For suppose D is such an H -bridge, let v_1, v_2 be its extremal attachments on P (these exist due to 5-connectivity), and let v be its sole attachment in H . Suffices now that we prove the following claims:

(4.7.B.1) *No member of $\mathcal{B}$ is adjacent to (v_1Pv_2) .*

(4.7.B.2) *Let $\mathcal{P}$ be the set of $((v_1Pv_2), H)$ -paths internally-disjoint of $D \cup P$. Then,*

(4.7.B.2.1) *$\mathcal{P}$ is nonempty;*

(4.7.B.2.2) *no member of $\mathcal{P}$ is adjacent to or meets a member of $\mathcal{B}$;*

(4.7.B.2.3) *the members of $\mathcal{P}$ have a common end in H ; and,*

(4.7.B.2.4) *there exists a member of $\mathcal{P}$ of order ≥ 3 .*

Indeed, assuming (4.7.B.1-2) (see proofs below), one may argue as follows. Let $u \in V(H)$ be the common end of the members of $\mathcal{P}$, existence of which follows from (4.7.B.2.3). By (4.7.B.2.2) and (4.7.B.2.4), there is an H -bridge C of B whose sole attachment in H is u ; and C meets vertices of $\mathcal{P}$. Let u_1 and u_2 be its extremal attachments on P . Clearly, (4.7.B.1-2) apply to C .

Since C meets $\mathcal{P}$, $u_1Pu_2 \cap v_1Pv_2 \neq \emptyset$. If $v_1Pv_2 \subseteq u_1Pu_2$, then there exists a $((u_1Pu_2), H)$ -path internally-disjoint of $D \cup C \cup P \cup \{u, v\}$ with an end disjoint of $\{u, v\}$. For otherwise since $|V(H)| \geq 3$ (as $Y \subseteq V(H)$ and stable) and by (4.7.B.1), $\{u, v, u_1, u_2\}$ is a 4-disconnector of G . Existence of such a path contradicts (4.7.B.2.3).

Consequently, $v_1Pv_2 \not\subseteq u_1Pu_2$ and (by symmetry) $u_1Pu_2 \not\subseteq v_1Pv_2$. Assume then, without loss of generality, that x, u_1, v_1, u_2, v_2, y appear in this order along P . In an analogous manner to the previous case, by (4.7.B.1), since $|V(H)| \geq 3$, and since $\{u, v, u_1, v_2\}$ is not a 4-disconnector of G , there is a $((u_1Pu_2), H)$ -path internally-disjoint of $D \cup C \cup P \cup \{u, v\}$ with an end disjoint of $\{u, v\}$. Since v_1Pv_2 and u_1Pu_2 overlap on P ; existence of such a path contradicts (4.7.B.2.3).

It remains to prove (4.7.B.1-2). To see (4.7.B.1), let P' be an induced xy -path obtained from P by replacing v_1Pv_2 with an induced v_1v_2 -path in $G[V(D) \cup \{v_1, v_2\}]$. If a member of $\mathcal{B}$ is adjacent to v_1Pv_2 , then P' contradicts (II).

By (4.7.B.1), since $|V(H)| \geq 3$, and since $\{v, v_1, v_2\}$ is not a 3-disconnector of G , $\mathcal{P}$ is nonempty. A member of $\mathcal{P}$ adjacent to or meeting a member of $\mathcal{B}$ implies that P' (as defined above) contradicts (4.7.b). Next, if $Q, Q' \in \mathcal{P}$ do not have a common end in H , then the union of Q, Q' , and v_1Pv_2 contains an H -ear. The path P' and the union of H and such an

ear contradict (I). Finally, if all members of $\mathcal{P}$ are edges, then by (4.7.B.1), (4.7.B.2.3), and since $|V(H)| \geq 3$, the set consisting of v, v_1, v_2 , and the common end of members of $\mathcal{P}$ is a 4-disconnector of G . $\square$

(4.7.C) $\mathcal{X}$ satisfies (4.7.e).

Proof. Let $B' \in \mathcal{B}$ and let $v \in V(B) \cap X$. If $V(B') \setminus \{v\} \neq \emptyset$, then the union of components of $B' - v$, namely $\mathcal{C}(B', v)$ is nonempty. Let $C \in \mathcal{C}(B', v)$ and let v_1, v_2 be the extremal attachments of C on P ; such vertices exist as $v \in N_G(C) \subseteq V(P) \cup \{v\}$ and $|N_G(C)| \geq 5$. Let P' be an induced xy -path obtained from P by replacing $v_1 P v_2$ with an induced $v_1 v_2$ -path in $G[V(C) \cup \{v_1, v_2\}]$.

H (which coincides with B , by (4.7.B)), is not adjacent to $(v_1 P v_2)$. (4.8)

Proof. Indeed, suppose to the contrary that H is incident to $(v_1 P v_2)$. Consider the following argument.

- (a) No member of $\mathcal{C}(B', v) \cup \mathcal{B}$ is adjacent to $(v_1 P v_2)$; for then P' is a path contradicting (4.7.b).
- (b) Since $\{v_1, v_2, v\}$ is not a 3-disconnector of G , the set $\mathcal{P}$ of $((v_1 P v_2), H)$ -paths internally-disjoint of $C \cup P$ is nonempty, by (a).
- (c) Clearly, no member of $\mathcal{P}$ is adjacent to or meets a member of $\mathcal{C}(B', v) \cup \mathcal{B}$, for then P' is a path contradicting (4.7.b).
- (d) Moreover, the members of $\mathcal{P}$ have a common end, say u , in H . Indeed, if not so and $Q, Q' \in \mathcal{P}$ have distinct ends in H , then the union of Q , Q' , and $(v_1 P v_2)$ contain an H -ear. The path P' and the union of H and such an ear contradict (I).
- (e) Since H coincides with B , the members of $\mathcal{P}$ are edges. It follows then that $\{v, u, v_1, v_2\}$ is a 4-disconnector of G . $\square$

To conclude our proof of (4.7.C), let $X' \subseteq X$ satisfying (i) $|X' \cap V(B')| = 1$ for every $B' \in \mathcal{B}$, and (ii) subject to (i) there is a $v \in X'$ such that $C(B, v) \neq \emptyset$ for some $B \in \mathcal{B}$.

Let $\mathcal{C} = \bigcup_{B' \in \mathcal{B}, v \in X' \cap V(B')} \mathcal{C}(B', v)$. Let $\mathcal{A}$ denote the subpaths of P each a union of extremal segments of overlapping and nested members of $\mathcal{C}$. By (4.8), H is not adjacent to $\text{int}A$ for any $A \in \mathcal{A}$. Let $A \in \mathcal{A}$ be maximal. The set comprised of X and the ends of A separates H from the members of $\mathcal{C}$ defining A . As, by assumption, $|X| < \kappa(G) - 2$, such a set has size $< \kappa(G)$. $\square$

■

As indicated above, in this paper, the counterpart of 4.7 is the following.

4.9. Suppose $\kappa(G) \geq 5$ and $H \cong K_{2,3} \subseteq G$ such that the 3-valent vertices of H are not adjacent in G . Then G contains an induced path P meeting no 2-valent vertex of H whose ends are the 3-valent vertices of H and with a P -bridge meeting at least two 2-valent vertices of H .

Proof. Let $V(H) = \{x_i\}_{i \in [5]}$ such that $d_H(x_1) = d_H(x_2) = 3$. Existence of disjoint $x_1 x_2$ -path and an $x_4 x_5$ -path in $G - x_3$, clearly implies the assertion. By the main result of [7, 8] (the

characterization of 2-linked graphs), nonexistence of such paths in $G - x_3$ occurs if and only if $G - x_3$ has an embedding in the plane such that x_1, x_4, x_2, x_5 appear on the outerface, say X (which is a circuit), in this clockwise order. As $\kappa(G - x_3) \geq 4$ and x_1, x_4, x_2, x_5 induce a C_4 , $X \cong C_4$.

Let C be an $\{x_1, x_2\}$ -circuit in $G - \{x_3, x_4, x_5\}$, and let P_1, P_2 be the two x_1x_2 -paths comprising C . Since $x_1x_2 \notin E(G)$, $\text{int}P_i \neq \emptyset$, $i = 1, 2$. The circuits C and X divide the plane into 4 regions:

$R_1 := \text{int}C_1$ with C_1 comprised of, say, P_1 and the edges $\{x_5x_1, x_5x_2\}$.

$R_2 := \text{int}C$ (not including C).

$R_3 := \text{int}C_2$ with C_2 comprised of, say, P_2 and the edges $\{x_4x_1, x_4x_2\}$.

$R_4 := \text{ext}X$.

Let $y \in N_G(x_3) \setminus \{x_1, x_2\}$ embedded in R_i , $i \in \{1, 2, 3\}$. Since $\kappa(G - x_3) \geq 4$, there is a $(y, \text{int}P_i)$ -path, Q , not meeting P_{3-i} for some $i = 1, 2$. Let $x \in V(C_i) \cap \{x_4, x_5\}$. There is an $(x, \text{int}P_i)$ -path, Q' , not meeting P_{3-i} . Since $Q \cup Q' \cup P_i \cup \{x_3\}$ contains an xx_3 -path disjoint of P_{3-i} the claim follows. ■

§4.2 Salami and Pie. The purpose of this section is to state the main result of Yu [9] that provides a characterization of the 4-connected plane graphs containing no TK_4 rooted at a prescribed set of 4 vertices.

Unless otherwise stated, throughout this section G is a 4-connected plane graph, and $W = \{w_1, w_2, w_3, w_4\} \subseteq V(G)$.

DEFINITION B.

A connected plane graph J is called a *configuration* if it has 6 distinct vertices $\{a_1, a_2, a_3, a_4, x, y\}$ such that:

(B.1) $A = \{a_1, a_2, a_3, a_4\}$ is stable and appears in this clockwise order on the outerwalk of J ,

(B.2) no disconnector T of J , $|T| \leq 3$, separates x or y from $A \setminus T$,

(B.3) no disconnector T of J , $|T| \leq 4$, separates $\{x, y\}$ from $A \setminus T$, and

(B.4) every disconnector T of J , $|T| \leq 3$, separates two vertices of $A \cup \{x, y\}$.

We refer to x and y as the *roots* of J and to the members of A as its *exits*.

Each graph in Figure 2 is a configuration with outerwalk, say C . The configurations (II)-(XI) each contains a 4-disconnector $S_x = \{s, s', t, t'\}$ separating $(A \setminus S_x) \cup \{y\}$ from x , and a 4-disconnector $S_y = \{p, p', q, q'\}$ separating $(A \setminus S_y) \cup \{x\}$ from y . The component of $J - S_x$ containing x is disjoint of the component of $J - S_y$ containing y . In configuration (I), if $x \notin V(C)$, then S_x as above exists. Otherwise, if $x \in V(C)$, we put $S_x = \{x\}$. A similar convention holds for y . We assume that $\{t, s, s', t'\}$ and $\{p, q, q', p'\}$ appear around x and y in these clockwise orders, respectively, as depicted in Figure 2.

In configuration (I), $a_1, t, s, p, q, a_2, a_3, a_4$ appear in this clockwise order on C ; and possibly $a_1 = t$ or $s = p$ or $q = a_2$. In configuration (II), $a_1, a_2, p, q, a_3, a_4, t, s$ appear in this clockwise order on C ; and possibly $s = a_1$ or $t = a_4$ or $p = a_2$ or $q = a_3$. In addition, s', p', q', t' lie on another face (other than C) and appear in this clockwise order on this face; possibly $s' = p'$ or $t' = q'$.

Configurations (III)-(XI) are variations of (II). In these configurations some constraints of (II) are relaxed. For instance, the presence of the face (other than C) containing s', p', q', t' is substituted by the constraint that $s' = p'$ or $t' = q'$. Also, it is no longer required that q and

t lie on C and so on. We refer and advise the reader to consult [9] for complete details.

It will be useful for us to observe the following.

4.10. *If J is a configuration as in Figure 2, then no exit of J is a common neighbor of its roots.*

Proof. An exit of J serving as a common neighbor of the roots of J contradicts (B.2). Let $x, y, A = \{a_i\}_{i=1,2,3,4}, S_x, S_y$, and C be as above. Let $a \in N(x) \cap N(y) \cap A$.

If $S_x \neq \{x\}$ and $S_y \neq \{y\}$, then $a \in A \cap S_x \cap S_y \neq \emptyset$; implying that $s' = p' \in A$ or $t' = q' \in A$. In each configuration, the latter translates into a disconnecter T , $|T| \leq 3$, of J separating at least one of x, y from $A \setminus T \neq \emptyset$; contradicting (B.2).

Suppose then that $S_x = \{x\}$; implying that J is configuration (I) and $x \in V(C)$. If $y \in V(C)$ as well, then at least one of the edges xa, ya forms a 2-disconnector T of J separating x or y from $A \setminus T \neq \emptyset$. Assume then that $S_y \neq \emptyset$. Consequently, $a_1 = a$, for otherwise $\{x, a\}$ is a 2-disconnector separating x_1 from y . Moreover, $a \in \{p', q'\}$. The latter implies that there is a disconnecter $T \subseteq \{p', q', q\}$, separating y (and x) from $A \setminus T$; contradicting (B.2). ■

DEFINITION C.

A pair (G, W) is called a *pie* if there is a facial circuit C of G , called the *mold*, and sets S_{w_i} , $i = 1, 2, 3, 4$, such that either:

(C.1) $w_i \in V(C)$ and then $S_{w_i} = \{w_i\}$, or

(C.2) G has a 4-disconnector S_{w_i} satisfying $|S_{w_i} \cap V(C)| = 2$, and separating w_i from $W \setminus \{w_i\}$. The components of $G - S_{w_i}$ containing w_i are disjoint and $w_j \notin S_{w_i}$ whenever $i \neq j$.

An illustration of a pie can be found in Figure 3(a). In the figure, the mold is drawn as the outerface.

DEFINITION D.

A pair (G, W) is called a *Salami separating* $\{w_1, w_2\}$ from $\{w_3, w_4\}$ if G contains distinct (not necessarily disjoint) 4-disconnectors T_i , $i \in [m]$ such that:

(D.1) each T_i separates $\{w_1, w_2\}$ from $\{w_3, w_4\}$,

(D.2) each T_i separates $\{w_1, w_2\}$ from $T_{i+1} \setminus T_i \neq \emptyset$,

(D.3) G has no 4-disconnector T separating $T_i \setminus T \neq \emptyset$ from $T_{i+1} \setminus T \neq \emptyset$,

(D.4) the 4-hammocks H and H' satisfying $\text{bnd}_G H = T_1$ and $\text{bnd}_G H' = T_m$ are minimal subject to containing $\{w_1, w_2\}$ and $\{w_3, w_4\}$, respectively.

(D.5) $J \in \{H, H'\}$ is a configuration as listed in Figure 2 such that there are bijections $\text{bnd}_G J \rightarrow A$ and $W \cap V(J) \rightarrow \{x, y\}$, where A and x, y are as in DEFINITION B.

An illustration of a Salami can be found in Figure 3(b). We refer to the hammocks H, H' of (D.4) as the *ends* of the Salami (G, W) .

A pair (G, W) , whether a pie or a Salami, has a set S_w associated with each $w \in W$. Such a set, if satisfies $S_w = \{w\}$, is called *elementary*; otherwise, S_w is the boundary of a 4-hammock containing w in its interior, and is then called *nonelementary*. A 4-hammock containing w and bounded by a nonelementary S_w has the members of S_w appear in a clockwise order along




Figure 2: Yu's configurations (copy of [9, Fig. 2]).




Figure 3: Yu's Salami and pie (copy of [9, Fig. 7]).

its outerwalk. We refer to this order as the *order* of S_w .

4.11. (Yu [9, Theorem 4.2])

G contains no TK_4 rooted at W if and only if (G, W) is a Salami or a pie.

In fact, Yu's argument for 4.11 [9, Theorem 4.2], which relies on [9, Theorems 2.1, 3.1], establishes the following more precise and useful fact.

4.12. If G contains no TK_4 rooted at W and there is a 4-disconnector T separating $\{w_1, w_2\}$ from $\{w_3, w_4\}$, then

(4.12.A) (G, W) is a Salami separating $\{w_1, w_2\}$ from $\{w_3, w_4\}$, and

(4.12.B) the ends of (G, W) are contained in distinct 4-hammocks whose boundary is T .

A consequence of 4.10 that will be useful for us is the following.

4.13. If three members of W have a common neighbor not in W , then (G, W) has a TK_4 rooted at W .

Proof. Suppose $c \in (\bigcap_{i=1}^3 N(w_i)) \setminus W$. By 4.10, (G, W) is not a Salami. Indeed, since any 4-disconnector separating two pairs of W contains c , both ends of (G, W) contain c as an exit; implying that at least one end has an exit that is a common neighbor of its roots, contradicting 4.10.

Suppose then that (G, W) is a pie and let C be its mold. Suppose $\{S_{w_1} \cap V(C), S_{w_2} \cap V(C), S_{w_3} \cap V(C), S_{w_4} \cap V(C)\}$ appear in this clockwise order on C . Since $\{w_1, c, w_3\}$ is not a 3-disconnector of G , at least one of w_1, w_3 is not on C ; so let S_{w_1} be nonelementary.

If $w_3 \notin V(C)$, then $c \in S_{w_1} \cap S_{w_3}$ and then G has a disconnector $T \subset (S_{w_1} \cap V(C)) \cup (S_{w_3} \cap V(C)) \cup \{c\}$, $|T| \leq 3$, separating w_2 from w_4 . On the other hand, if $w_3 \in V(C)$ then G has a disconnector $T \subset (S_{w_1} \cap V(C)) \cup \{c, w_3\}$, $|T| \leq 3$, separating w_2 from w_4 . ■

We shall use the following observation that is a consequence of **4.13**.

4.14. *If $H \cong K_{2,3} \subseteq G$ and G is a nonplanar 5-connected apex graph, then either $TK_5 \subseteq G$, or no 3-valent vertex of H is an apex vertex of G .*

Proof. Fix an embedding of G in the plane and treat G as a plane graph. Let $u \in V(H)$ be 3-valent and let $W \subset N_G(u)$ consist of the three 2-valent vertices of H and a fourth vertex not in $V(H)$. Since 3 members of W have a common neighbor not in W , it follows, by **4.13**, that if u is an apex vertex of G , then $G-u$ contains a TK_4 rooted at W ; implying that $TK_5 \subseteq G$. ■

The facial circuits of a 4-connected planar graph are its induced nonseparating circuits and are unique [6]. Let $C \subset G$ be an induced 4-circuit with $V(C) = \{x_1, x_2, x_3, x_4\}$ such that $x_1 x_3 \notin E(G)$. Let H be a 4-hammock of G such that $x_1 \in \text{int}_G H$ and $x_3 \notin V(H)$. Clearly, $x_2, x_4 \in \text{bnd}_G H = \{u_0, u_1, u_2, u_3\}$, where the latter appear in this clockwise order along the outerwalk of H (here we refer to the embedding of H induced by the embedding of G). It is not necessarily true that C is a facial circuit of G . However,

$$\{x_2, x_4\} = \{u_{i(\text{mod } 4)}, u_{i+1(\text{mod } 4)}\}, 1 \leq i \leq 4, \text{ if and only if } C \text{ is a facial circuit of } G. \quad (4.15)$$

To see (4.15), suppose that $\{x_2, x_4\} = \{u_{i(\text{mod } 4)}, u_{i+1(\text{mod } 4)}\}$, $i = \{1, 2, 3, 4\}$ and C is a separating circuit of G . That is, at least one of $H \cap \text{int} C$, $(G-H) \cap \text{int} C$ is a nonempty graph. By the assumption on $\{x_2, x_4\}$ and since H is a hammock, at least one of $\{x_1, x_2, x_4\}$, $\{x_2, x_3, x_4\}$ is a 3-disconnector of G ; a contradiction. The converse is clearly true.

Suppose, next, that J is a configuration as in Figure 2 with x_1 and x_3 as its roots such that $x = x_1$ and $y = x_3$, say, and S_x, S_y are both nonelementary. Clearly, $x_2, x_4 \in S_x \cap S_y$. If (B.2-4) are to be satisfied subject to the allowed vertex identifications specified for configurations in Figure 2, then

(M.1) $\{x_2, x_4\} = \{s, s'\} = \{p, p'\}$ in configuration (I), or

(M.2) $\{x_2, x_4\} = \{s', t'\} = \{p', q'\}$ in configurations (II)-(XI).

A similar situation arises when only one of S_x and S_y is nonelementary and also when S_x and S_y are part of a pie and not a configuration. Consequently, we observe the following.

4.16. *Let x, y be part of an induced 4-circuit such that x and y are nonadjacent. Let $z \in \{x, y\}$.*

(4.16.A) *If J is a configuration as in Figure 2 with roots x, y , then an S_z that is nonelementary satisfies $V(C) \setminus \{x, y\} \subset S_z$ and the members of $V(C) \cap S_z$ appear as in (M.1) and (M.2).*

(4.16.B) *If (G, W) is a pie with mold F , where $W = \{x, y, \ell, \ell'\}$ and $|V(C) \cap W| = 2$, then $\{S_x \cap F, S_y \cap F, S_\ell \cap F, S_{\ell'} \cap F\}$ appear in this clockwise order along F ; an S_z that is nonelementary satisfies $V(C) \setminus \{x, y\} \subset S_z$ such that the members of $V(C) \cap S_z$ appear consecutively in the order of S_z : one on F and the other not on F .*

The following is now an exercise.

4.17. *Suppose w_1, w_2 are a part of an induced 4-circuit C of G such that $w_1 w_2 \notin E(G)$ and $|V(C) \cap W| = 2$. Then, G has a TK_4 rooted at W or C is a facial circuit of G .*

§4.3 Proof of 4.3. Throughout this section, G is a 5-connected nonplanar apex graph satisfying $TK_5 \not\subseteq G$; K is as in (3.1); a is an apex vertex of G as in 4.3; and P is an induced x_1x_2 -path satisfying (4.2).

By 1.1 and 2.3, to prove 4.3, suffices that we prove the following claims:

(4.3.A) *If $a \in \{x_3, x_4, x_5\}$, then $K_4^- \subseteq G$ or G has a planar hammock or $TK_5 \subseteq G$.*

(4.3.B) *If $a \notin V(K)$, then $K_4^- \subseteq G$ or $G - a$ is nonplanar.*

Claim (4.3.B) is an exercise; though not necessarily a short one. Nevertheless, it becomes a routine case analysis due to (4.2) (which is a consequence of 4.7). We concentrate on (4.3.A) only.

§4.3.1 Proof of (4.3.A). Throughout this proof $a = x_3$ and C is the C_4 induced by $V(K) \setminus \{a\}$. The facial circuits of a 4-connected planar graph are its induced nonseparating circuits and are unique [6]. Consequently, by (4.1) and 4.17,

$$G - P \text{ is a chain, } a \text{ is a cut vertex of } G - P, \text{ and } C \text{ is a facial circuit of } G - a. \quad (4.18)$$

To see (4.18), note that if $G - P - a$ contains an x_4x_5 -path P' , then, in $G - a$, the set $\{x_1, x_2, x_4, x_5\}$ forms an induced 4-circuit separating $\text{int}P$ from $\text{int}P'$ each is nonempty by (4.1). Therefore, by 4.17 applied to $(G - a, \{x_1, x_2, y, y'\})$, $y, y' \in N_G(a) \setminus V(K)$ (these exist by (4.1)), it follows that $TK_5 \subseteq G$. This establishes that $G - P$ is a chain and that a is its cut vertex. Consequently, $G - a$ has an embedding in which P and $G - P - a$ are both embedded in $\text{int}C$ or $\text{ext}C$ and thus C is a facial circuit of $G - a$.

OUTLINE. We now outline the reminder of our proof for (4.3.A); adjourning technical details until later parts of this section. Throughout, the remainder of this section W is a set of the form $\{x_1, x_2, \ell, \ell'\}$, where $\ell, \ell' \in N_G(a) \setminus V(K)$, $G - a$ is embedded in the plane such that C is its outer face; such an embedding is called II. We begin by proving (see proof below) that

(4.3.A.1) *there exist $\ell, \ell' \in N_G(a) \setminus V(K)$ such that $(G - a, \{x_1, x_2, \ell, \ell'\})$ is a Salami.*

We distinguish between two cases:

CASE A. Either $G - a$ has a 4-disconnector T separating $\{x_1, \ell\}$ from $\{x_2, \ell'\}$ for some $\ell, \ell' \in N_G(a) \setminus V(K)$; or

CASE B. T as above does not exist in $G - a$ for any $\ell, \ell' \in N_G(a) \setminus V(K)$.

Our argument for CASE A is as follows. Let $\mathbf{L}$ be the Salamies $(G - a, W)$ separating $\{x_1, \ell\}$ from $\{x_2, \ell'\}$ for some $\ell, \ell' \in N_G(a) \setminus V(K)$. The assumption of this case that T exists and 4.12 imply that $\mathbf{L} \neq \emptyset$. For $(G - a, W) \in \mathbf{L}$, let $\alpha(G - a, W) = |V(J) \cup V(J')|$, where J and J' are the ends of $(G - a, W)$.

$$\text{Choose } (G - a, W) \in \mathbf{L} \text{ minimizing } \alpha(G - a, W). \quad (4.19)$$

SETTING A. For $(G - a, W)$ as in (4.19), let J_i denote the end of $(G - a, W)$ containing x_i , $i = 1, 2$. Put $x \in \{x_1, x_2\}$ and $y \in \{\ell, \ell'\}$. Let $J \in \{J_1, J_2\}$ such that $x, y \in \text{int}_{G-a} J$. Let $S_x, S_y, A = \{a_1, a_2, a_3, a_4\} = \text{bnd}_{G-a} J$ be as these are defined in §4.2 and Figure 2. Since any 4-disconnector separating x_1 and x_2 contains $\{x_4, x_5\}$, we have that $\{x_4, x_5\} \subseteq \text{bnd}_{G-a} J$.

A proof of (4.3.A) then follows from the following two claims (that are proven below):

(4.3.A.2) $J - \{x_4, x_5\}$ contains two paths Y and Q such that:

(4.3.A.2.1) Y is an xa_2 -path and Q is an xa_3 -path; and,

(4.3.A.2.2) $V(Y) \cap V(Q) = \{x, y\}$.

(4.3.A.3) $J - \{x_4, x_5\}$ contains an $(x, \{a_2, a_3\})$ -2-fan with one of its members containing y .

Proof that (4.3.A.2-3) imply (4.3.A). Let $\text{bnd}_{G-a}J_1 = U = \{u_1, u_2, u_3, u_4\}$ and let $\text{bnd}_{G-a}J_2 = Z = \{z_1, z_2, z_3, z_4\}$ such that $u_1 = x_4 = z_1$, $u_4 = x_5 = z_4$, $\{u_1, u_2, u_3, u_4\}$ and $\{z_1, z_2, z_3, z_4\}$ appear in these clockwise order on the outerwalks of J_1 and J_2 , respectively, with respect to Π .

Let $Y, Q \subseteq J_1$ be as in (4.3.A.2), where $x = x_1, y = \ell$, $a_2 = u_2$, and $a_3 = u_3$. Let $F \subseteq J_2$ be as in (4.3.A.3) where $x = x_2, y = \ell'$, $a_2 = z_2$, and $a_3 = z_3$. Let Y' be the member of F containing ℓ such that, without loss of generality, $z_2 \in V(Y')$. Let Q' be the other member of F . Let $F' = \{Y'', Q''\}$ be a $(\{u_2, u_3\}, \{z_2, z_3\})$ -2-linkage in $G - \text{int}_{G-a}J_1 - \text{int}_{G-a}J_2 - \{a, x_4, x_5\}$. Such a linkage clearly exists as $\kappa(G - \{a, x_4, x_5\}) \geq 2$. By planarity, we may assume that Y'' is a u_2z_2 -path and that Q'' is a u_3z_3 -path. Let X_Y and X_Q be the x_1x_2 -paths in $G - a$ that are the union of Y, Y', Y'' and Q, Q', Q'' , respectively. Let F'' be an (x_4, X_Y) -4-fan in $G - a$ with two of its members the edges $\{x_1x_4, x_2x_4\}$ and minimizing the length of its members. Let X, X' be the remaining members of F'' . Such do not meet X_Q , by planarity.

Now, the path X_Q satisfies (F.2); implying that $TK_5 \subseteq G$. To see this, suffices to show that $G - X_Q - a + \ell$ contains an $\ell\ell'$ -path containing x_4 (recall that $x_3\ell, x_3\ell' \in E(G)$). Such a path clearly exists in $X \cup X' \cup (X_Y - \{x_1, x_2\})$. ■

With CASE A resolved, we may now assume, due to 4.12, that in $G - a$

there are no $\ell, \ell' \in N_G(a) \setminus V(K)$ and a 4-disconnector separating $\{x_1, \ell\}, \{x_2, \ell'\}$. (4.20)

Consequently, by (4.3.A.1), there are $\ell, \ell' \in N_G(a) \setminus V(K)$ such that $(G - a, W)$ is a Salami separating $\{x_1, x_2\}$ from $\{\ell, \ell'\}$. This and (4.20) clearly imply that

(4.3.A.4) *there exist $\ell, \ell' \in N_G(a) \setminus V(K)$ such that $(G - a, W)$ is a Salami such that*

(4.3.A.4.1) x_1, x_2 are contained in a common end, and

(4.3.A.4.2) *for $i = 1, 2$, S_{x_i} , if nonelementary, defines a 4-hammock H_i of $G - a$ satisfying*

$$x_i \in \text{int}_{G-a}H_i, \text{bnd}_{G-a}H_i = S_{x_i}, \text{ and } N_G(a) \cap \text{int}_{G-a}H_i = \{x_i\}.$$

Proof that (4.3.A.4) implies (4.3.A). Let J be the end of $(G - a, W)$ containing $\{x_1, x_2\}$. Clearly, a nonelementary S_{x_i} satisfies $\{x_4, x_5\} \subset S_{x_i}$. We prove the following claims (a) and (b).

(a) S_{x_i} is nonelementary for at least one $i = 1, 2$.

Proof. Otherwise, the end J of $(G - a, W)$ containing x_1, x_2 is configuration (I) from Figure 2 with both x_1, x_2 appearing on its outerwalk. Since $x_1x_2 \notin E(G)$, by (4.1), at least one of x_4, x_5 also lies on the outerwalk of J or $\{x_1, x_2, x_4, x_5\}$ contains a 3-disconnector of $G - a$. As $\{x_1, x_2, x_4, x_5\}$ define an induced C_4 , at least one of $\{x_4, x_5\}$ has degree at most 2 in $G - a$ which is a contradiction. □

(b) If H_i is as in (4.3.A.4.2) for some $i = 1, 2$, then H_i is a trivial 5-hammock of G so that $N_G(x_i) = \text{bnd}_{G-a}H_i \cup \{a\}$.

Proof. Since $N_G(a) \cap \text{int}_J H_i = \{x_i\}$, by (4.3.A.4.2), then $\text{bnd}_G H_i = \text{bnd}_{G-a} H_i \cup \{x_i\} = S_{x_i} \cup \{x_i\}$. As $|V(G)| > |V(H_i)|$, we may assume that H_i is not a planar hammock or we are done; and thus, $|V(H_i)| \leq 6$. If $|V(H_i)| = 6$, then the (sole) vertex v in $\text{int}_G H_i$ has $N_G(v) = \text{bnd}_G H_i$; implying that $\{v, x_1, x_4, x_5\}$ induce a K_4^- . As H_i is a 4-hammock of $G - a$, it follows that H_i is a trivial 5-hammock of G and that $N_G(x_i) = \text{bnd}_{G-a} H_i \cup \{a\}$. $\square$

To conclude, let then S_{x_1} be nonelementary and H_1 (as in (4.3.A.4.2)) be a trivial 5-hammock of G , by (a) and (b) (above). We may assume that S_{x_2} is nonelementary. If not so, then J is configuration (I) from Figure 2 with x_2 appearing on its outerwalk. Since C is a 4-face of $G - a$, then by 4.16, triviality of H_1 , (B.2-4), and 4-connectivity of $G - a$, at least one of x_4, x_5 also lies on the outerwalk of J and consequently (by planarity of $G - a$) has degree at most 3 in $G - a$; which is a contradiction.

By (4.3.A.4.2) and (b) (above), S_{x_2} being nonelementary, implies that there is an H_2 (as in (4.3.A.4.2)) that is a trivial 5-hammock of G . Since C is a face, J is configuration (I)-(V). If J is configuration (I), then at least one of x_4, x_5 lies on the outerwalk of J , by 4.16. Triviality of H_1, H_2 , and since a is not adjacent to $\{x_4, x_5\}$, imply that such a vertex has degree ≤ 4 in G ; a contradiction.

Suppose then that J is configuration (II)-(V). Let $S_{x_1} = \{s, s', t, t'\}$, $S_{x_2} = \{p, p', q, q'\}$ such that $\{s, s', t, t', p, p', q, q'\}$ are as in §4.2 and Figure 2. Triviality of H_1, H_2 , observation 4.16, the fact that $N_G(x_j) \subseteq V(J)$, for $j = 4, 5$, and since $N_G(x_i) = S_{x_i} \cup \{a\}$, $i = 1, 2$, imply that J has a 3-disconnector of the form $\{s, p, v\}$ or $\{t, q, v\}$, where $v \in \{x_4, x_5\}$, contradicting at least one of (B.2-4). $\blacksquare$

PROOFS. To conclude our proof of (4.3.A), it remains to prove claims (4.3.A.1-3). This is done next.

Proof of (4.3.A.1). By the assumption that $\kappa(G - P) = 1$ and (4.18), $G - P$ is the union of an $x_4 a$ -subchain X_1 and an $x_5 a$ -subchain X_2 such that $V(X_1) \cap V(X_2) = \{a\}$. Let $y_i \in N_G(a) \cap V(X_i)$, $i = 1, 2$. Let P_1 be an $x_4 y_1$ -path in X_1 and let P_2 be an $x_5 y_2$ -path in X_2 . Clearly, P_1 , and P_2 are disjoint and $V(P) \cap V(P_i) = \{x_i\}$, $i = 1, 2$.

Assume, towards contradiction, that $(G - a, \{x_1, x_2, y_1, y_2\})$ is a pie and let F be its mold. Since C is a 4-face and $\kappa(G - a) \geq 4$, we may assume that $\{S_{x_1} \cap F, S_{x_2} \cap F, S_{y_1} \cap F, S_{y_2} \cap F\}$ appear in this clockwise order along F . Clearly, S_{x_i} , $i = 1, 2$, if nonelementary, contains x_4 and x_5 and these appear as specified in (4.16.B). Thus, since $x_4, x_5 \notin V(P)$, we have that $(V(P) \cap V(P_i)) \setminus \{x_i\} \neq \emptyset$ for at least one $i = 1, 2$; contradiction. $\blacksquare$

For the proofs of (4.3.A.2-3), we require some preparations. Throughout the reminder of this section, we refer to the notation and terminology of SETTING A above.

Any 4-disconnector separating x_1 from x_2 contains x_4 and x_5 ; hence, $x_4, x_5 \in A$. Consequently, S_x , if nonelementary, satisfies $\{x_4, x_5\} \subset S_x \cap A$. This cannot occur in configuration (I) without contradicting (B.2). On the other hand, if S_x is elementary and (thus) J is configuration (I), then x is on the outerwalk of J and is adjacent to two members of A . This also cannot occur in configuration (I) without contradicting (B.2). We infer that:

$$J \text{ is not configuration (I).} \tag{4.21}$$

Since x_1, x_2 are on the outer face of G (recall Π), we may assume that $a_1 = x_4 = s, a_4 = x_5 = t$, $S_x = \{s, s', t, t'\}$, and $S_y = \{p, p', q, q'\}$, where $\{s, s', t, t', p, p', q, q'\}$ are as in §4.2 and Figure 2. Let H_x be the 4-hammock of J satisfying $x \in \text{int}_J H_x$ and $\text{bnd}_J H_x = S_x$. In a similar manner define H_y for y . Next we prove claims (4.3.A.5-7).

(4.3.A.5) $V(H_x) = S_x \cup \{x\}$ so that $N_{G-a}(x) = S_x$.

Proof. By the minimality of J (see (4.19)) and 4.12, $N_G(a) \cap \text{int} H_x = \{x\}$. Therefore, $\text{bnd}_G H_x = S_x \cup \{x\}$. As we may assume that H_x is not a planar hammock of G , it follows that $|V(H_x)| \leq 6$. If equality holds, then there exists a (single) vertex $v \in \text{int}_G H_x$ with $N_G(y) = \text{bnd}_G H_x = S_x \cup \{x\}$; and then $\{x, v, x_4, x_5\}$ induce a K_4^- . ■

(4.3.A.6) $S_x \cap N_G(a) = \emptyset$.

Proof. As $\{x_4, x_5\} \subset S_x \cap \text{bnd}_{G-a} J$, suffices that we show that $s', t' \notin N_G(a)$. Indeed, assume $s' \in N_G(a)$. Then $(G - a, \{x, s', \{x_1, x_2\} \setminus \{x\}, \{\ell, \ell'\} \setminus \{y\}\})$ is a Salami with $\{x, s'\}$ at a common end J' , by 4.12. The 4-hammock J' is not configuration (I), by (4.21). Thus, J' has a 4-disconnector separating x from s' , which is a contradiction as $xs' \in E(G)$, by (4.3.A.5). ■

A consequence of (4.3.A.5-6) is that

$$d_J(s'), d_J(t') \geq 5. \quad (4.22)$$

(4.3.A.7) *There are disjoint paths P' and Q' satisfying the following:*

(4.3.A.7.1) *P' is an $s'p'$ -path and Q' is a $t'q'$ -path;*

(4.3.A.7.2) *neither meets $A \cup \{p, q, x\} \cup (V(H_y) \setminus \{p', q'\})$ (so that $V(P') \cup V(Q') \subseteq J$).*

(4.3.A.7.3) *Moreover, if P' and Q' are not part of the boundary of a face of J containing $\{s', t', p', q'\}$, then at least one of them consists of a single vertex.*

Proof. Since J is not configuration (I), by (4.21), we have that either $\{s', t'\} \cap \{p', q'\} = \emptyset$, and then $\{s', t', p', q'\}$ are cofacial; or $\{s', t'\} \cap \{p', q'\} \neq \emptyset$. We consider these two cases.

Suppose that there is a face f containing $\{s', t', p', q'\}$. If $V(f) \setminus (V(H_y) \cup V(H_x))$ meets $A \cup \{p, q\}$, then the degree of s' or t' is < 5 ; or there is a disconnector T , $|T| \leq 3$ separating y from some member of A , or there is a disconnector of $G - a$ of size ≤ 3 . These contradict (4.22), (B.2), and 4-connectivity of $G - a$, respectively. The paths required then lie on the boundary of f ; and (4.3.A.7.1) is satisfied.

If a face as above does not exist, then $|\{s', t'\} \cap \{p', q'\}| \geq 1$. By (4.21), we may assume then that $p = a_2$ (the case that $q = a_3$ is symmetrical), and therefore $s' \neq p'$, or $\{x, s', p\}$ separate y from a_1 in J ; contradicting (B.2). Consequently, $t' = q'$, and one of the required paths is a single vertex. A path as P' is then contained in an $(s', \{a_1, p = a_2, t' = q', p'\})$ -4-fan which exists in $G - a$, contained in J , and does not meet x , by (4.3.A.5). ■

We are now ready to prove claims (4.3.A.2-3).

Proof of (4.3.A.2). Let F be a (y, S_y) -4-fan; clearly, $F \subseteq H_y$. Let P' and Q' be an $s'p'$ -path and a $t'q'$ -paths, respectively, as in (4.3.A.7). If a_2 and a_3 coincide with p' and q' , respectively, then it is easy to see that Y and Q are contained in $F \cup P' \cup Q' \cup \{xs', xt'\}$, respectively. Therefore, let us assume that $a_3 \neq q$ (the argument for a_2 is symmetrical) which implies that p lies on the outerwalk of J (by the list of possible configurations).

Let us show that there is an $r \in N_J(a_3) \setminus V(Q')$. For suppose that $N_J(a_3) \subseteq V(Q')$. Let, then, $b \in N_J(a_3)$ such that $t'Q'b$ is minimal. If P' and Q' are parts of the boundary of a face containing $\{s', t', p', q'\}$, then $\{t', b, a_4\}$ is a 3-disconnector of J contradicting (B.2). To see this, note that $d_J(t') \geq 5$, by (4.22), and that $N_{G-a}(x) = S_x$. Minimality of b then implies the assertion.

Consequently, at least one of P' and Q' is a single vertex. Since p is on the outerwalk of J , $s' = p'$ implies that $\{x, s' = p', p\}$ is a 3-disconnector of J contradicting (B.2). Thus, $t' = q' = b$ and then $\{x, t' = b = q', a_3\}$ is a 3-disconnector of J contradicting (B.2).

We have shown that r as above exists. Since p is on the outerwalk of J , then $Q' \cup \{x, a_4, a_3, p\}$ separates r from a_4 . Also, observe that $r \neq x$, by (4.3.A.5), and that $r \notin \text{int}_J H_y$.

Let F' be as follows: if q is not on the outerwalk of J , then F' is an $(r, \{q', q, a_4, a_2\})$ -4-fan. Otherwise, F' is an $(r, \{q', q, a_3, a_2\})$ -4-fan. Let Q_q and $Q_{q'}$ be the members of F' ending at q and q' , respectively.

(i) By planarity and disjointness of Q_q and $Q_{q'}$, $V(Q_{q'}) \cap V(Q') = \emptyset$.

(ii) By definition of F' and planarity, $Q_{q'} \subseteq J - \text{int}_J H_y$.

To state Y and Q properly, define r to be a_3 if $a_3 = q$, and $r \in N_J(a_3) \setminus V(Q')$, otherwise. In an analogous manner define r' for a_2 with p replacing q . This way we may define a path P_p as a pa_2 -path disjoint of P' and contained in $J - \text{int}_J H_y$ in a symmetrical construction to that yielding Q_q (above). Planarity asserts that we may choose P_p and Q_q disjoint. Clearly, Y, Q as required are contained in $F \cup P' \cup Q' \cup P_p \cup Q_q \cup \{xs', xt', ra_3, r'a_2\}$. ■

Proof of (4.3.A.3). Let P', Q' be as in the proof of (4.3.A.2). Constructing an xa_2 -path (resp., xa_3 -path) containing P' (resp., Q') and y can be done precisely as in the proof of (4.3.2). We concentrate on constructing an xa_3 -path not meeting y so that together these paths would constitute the required fan.

REMARK. In the sole case that $q = a_3$ and p not on the outerwalk of J we shall prefer to construct the paths so that the xa_3 -path meets y ; the construction of the other path in this case is symmetrical to what follows.

Let $r \in N_J(a_3) \setminus \text{int}_J H_y$; such a vertex clearly exists if $a_3 \neq q$. Nonexistence of such a vertex in case $a_3 = q$, implies that $\{p, p', q'\}$ is a 3-disconnector of J separating y from $\{a_1, a_4\}$, contradicting (B.2).

Let F' be an $(r, \{q', q, a_3, a_4\})$ -4-fan. By planarity and the disjointness of the members of F' , the path $P_{q'} \in F'$ ending at q' is contained in $J - \text{int}_J H_y$, and the required remaining xa_3 -path is contained in $Q' \cup P_{q'} \cup \{xt', ra_3\}$. ■

§4.4 Proof of 4.5. The reader should be reminded of the AGREEMENT specified in §2. Throughout this section, G is a 5-connected nonplanar apex graph satisfying $TK_5 \not\subseteq G$; P is an induced x_1x_2 -path satisfying (4.4), where $\{x_i\}_{i=1}^k$ and K are as in (3.1); H is as in (4.4); and a is an apex vertex of G as in 4.5.

Claim 4.5 follows from the following two claims; of which (4.5.A) is an exercise.

(4.5.A) $a \in \{x_4, x_5\}$; otherwise $TK_5 \subseteq G$ or $G - a$ is nonplanar.

(4.5.B) If $a \in \{x_4, x_5\}$, then $TK_5 \subseteq G$ or G contains a planar hammock or $K_4^- \subseteq G$.

COMMON THEME. The following is a common theme of the proofs of (4.5.A-B): Let $x \in \text{int}P$ and let $y, z \in N_H(x)$. A $\{yz, x_4, x_5\}$ -circuit in $H + yz$ clearly implies an $\{x, x_4, x_5\}$ -circuit in $H + x$ satisfying (F.2). Let then $\mathcal{X}$ be a $\{yz, x_4, x_5\}$ -frame in $H + yz$, which exists by 3.4.

Let $H_{y,z}, H_{x_4}, H_{x_5}$ be the seeded $\mathcal{X}$ -hammocks containing $\{y, z\}, x_4, x_5$, respectively. Let F_1 and F_2 denote the components of such a frame.

Proof of (4.5.A). It is not hard to see that $a \in V(P) \cup \{x_3\}$ contradicts planarity of $G - a$ or 5-connectivity of G . Suppose then that $a \in V(H)$. Choose x, y, z as in the COMMON THEME; below we refer to the notation and terminology set in the COMMON THEME. In $H - yz$ there is an $\{x_4, x_5\}$ -circuit meeting the ends of $H_{y,z}$ and not meeting $\text{int}_H H_{y,z}$. Consequently, $H - a - yz$ contains paths P' and Q satisfying:

- (i) P' is an $x_4 x_5$ -path,
- (ii) Q is a ww' -path such that $w \in N_{H_{y,z}}(x)$, $w' \in \text{int} P'$,
- (iii) $V(Q) \cap V(P') = \{w'\}$. Thus, unless $a \in \{x_4, x_5\}$, then a $TK_{3,3}$ is contained in the union of: P', Q, P , and $V(K) \setminus \{x_3\}$. ■

Proof of (4.5.B). Assume, without loss of generality, that $a = x_5$, and choose x, y, z as in the COMMON THEME; below we refer to the notation and terminology set in the COMMON THEME. We argue as follows: (a) $x_4 \in \text{int}_H H_{x_4}$, by definition of frames, so that $|V(H_{x_4})| \geq 3$.

(b) By 5-connectivity, H_{x_4} , having order ≥ 3 , has at least two attachments on P . Let Q denote the extremal segment of H_{x_4} on P ; such does not consist of a single edge for then the ends of Q and the ends of H_{x_4} form a 4-disconnector of G .

(c) $N_G(x_5) \cap \text{int} Q = \emptyset$. For suppose $\ell \in N_G(x_5) \cap \text{int} Q$. Then, due to planarity, there exists an $\ell' \in N_{H_{x_4}}(\ell)$. An $\ell' x_5$ -path containing x_4 clearly exists in H . Indeed, a seeded- ℓ' -path in H_{x_4} and a seeded-through path in H_{x_5} are easily extended into such a path. Consequently, an $\{\ell, x_4, x_5\}$ -circuit satisfying (F.2) exists in $H + \ell$.

(d) Since the ends of Q and the ends of H_{x_4} do not form a 4-disconnector of G , x_3 is adjacent to $\text{int} Q$. We may assume that x_3 , the ends of Q , and the ends of H_{x_4} do not form the boundary of a planar hammock. Since $x_4 \in \text{int}_H H_{x_4}$, then these form the boundary of a 5-hammock of G of order 6 with x_4 as the sole vertex in the interior of this hammock. This implies that $x_4 x_3 \in E(G)$; contradicting (4.1). ■

§4.5 Proof of 4.6. Throughout this section, G is a 5-connected nonplanar apex graph satisfying $TK_5 \not\subseteq G$; P is an induced $x_1 x_2$ -path satisfying (4.4), where $\{x_i\}_{i=1}^k$ and K are as in (3.1); H is as in (4.4); and a is an apex vertex of G as in 4.6.

Claim 4.6 follows from the claims (4.6.A-B). Tough not necessarily short, (4.6.A) is an exercise which we omit.

(4.6.A) *If $a \notin \{x_4, x_5\}$, then $G - a$ is nonplanar.*

(4.6.B) *If $a \in \{x_4, x_5\}$, then $TK_5 \subseteq G$ or $K_4^- \subseteq G$ or G has a planar hammock.*

Proof of (4.6.B). Assume, without loss of generality, that $a = x_5$. The circuit $C = P + x_1 x_4 + x_2 x_4$ separates x_3 from $H - a$. Fix an embedding of $G - a$ such that x_3 lies in $\text{ext} C$ and $H - a$ lies in $\text{int} C$. Let B be the leaf block of H containing x_4 . Let H' be the subchain of H containing all blocks of H other than B . Let y denote the cut vertex of H present in B ; by 4.5, H' and y exist.

OUTLINE. We now outline our argument for proving 4.6; adjourning technical details to the end of the section. Proofs of claims hereafter follow below. Initially we prove that

(4.6.B.1) B is nontrivial (i.e., does not consist of a single edge).

By (4.6.B.1), the boundary of the outer face of B is a circuit comprised of two x_4y -paths. An attachment vertex of B in $\text{int}P$ is incident to at least one of these paths. Planarity asserts that there is at most one vertex in $\text{int}P$ that is adjacent to both of these paths. Consequently, there are attachments $z, z' \in \text{int}P$ of B such that zPx_2 contains all attachments adjacent to one of these x_4y -paths, and $z'Px_1$ contains all attachments of B adjacent to the other path. These segments of P are disjoint, unless $z = z'$. We will prove that

(4.6.B.2) $N_G(x_5) \cap (z'Px_1) = N_G(x_5) \cap (zPx_2) = \emptyset$.

(4.6.B.3) $|\{z, z', x_1, x_2\}| = 4$.

(4.6.B.4) $N_G(x_3) \cap (z'Px_1) \neq \emptyset$ or $N_G(x_3) \cap (zPx_2) \neq \emptyset$.

(4.6.B.5) H' does not consist of a single edge (i.e. x_5y).

Let $w, w' \in V(P)$ be attachments of $H' - x_5$ on P such that wPx_2 is minimal and $w'Px_1$ is minimal; these exist by (4.6.B.5). By planarity of $G - a$, $w, w' \in zPz'$. We will show that

(4.6.B.6) $N_G(x_3) \cap (wPw') \neq \emptyset$.

Let $h_1 \in N_G(x_3) \cap (wPw')$, by (4.6.B.6). Let $h'_1 \in N_G(h_1) \cap V(H' - x_5)$. Let $h_2 \in N_G(x_3) \cap (z'Px_1)$, without loss of generality, by (4.6.B.4). By (4.6.B.2), h_2 is at least 5-valent in $G - x_5$. Planarity then implies that there is an $h'_2 \in N_G(h_2) \cap V(B) \setminus \{x_4, y\}$. Indeed, if $h_2x_4 \in E(G)$, then since P is induced x_1 is at most 4-valent in G ; contradicting 5-connectivity. Let $b \in N_G(x_2) \cap V(B) \setminus \{y, x_4\}$. Such a vertex exists for otherwise $x_2y \in E(G)$, implying that $z = x_2$; contradicting (4.6.B.3). We will see that

(4.6.B.7) $B - x_4 - y$ contains a bh'_2 -path.

With the above claims in place, we construct a TK_5 in G whose branch vertices are x_3, h_1, h_2, x_1, x_2 as follows. Since $\{h_1, h_2, x_1, x_2\} \subset N_G(x_3)$, suffices that we show that $G - x_3$ has a TK_4 rooted at $\{h_1, h_2, x_1, x_2\}$. Since C contains all these vertices, such a rooted TK_4 is implied by existence of an x_1h_1 -path Q_1 and an x_2h_2 -path Q_1 in $G - x_3$ that are internally-disjoint of C and disjoint from one another. Put $Q_1 = x_1x_5 + x_5h_3 + Q'_1 + h_1h'_1$, where $h_3 \in N_{H'}(x_5)$ and Q'_1 is an h'_1h_3 -path in $H' - x_5$. The path Q'_1 exists since is not a cut vertex of H' as it is contained internally in one of its leaf blocks. Next, put $Q_2 = x_2b + Q'_2 + h'_2h_2$, where Q'_2 is as in (4.6.B.7).

PROOFS. It remains to prove (4.6.B.1-7).

Proof of (4.6.B.1). If B is trivial, then it consists of the edge x_4y , as x_4 is contained internally in B . Planarity, P being induced, the P -bridge meeting x_3 is a singleton, imply that there is a $w \in \{x_1, x_2\}$ and a vertex in $\text{int}P$ such that these two vertices and x_4 define a triangle. Planarity then implies that $wy \notin E(G)$ and thus w is at most 4-valent in G ; contradicting 5-connectivity. $\square$

Proof of (4.6.B.2). Suppose, to the contrary, that $r \in N_G(x_5) \cap (zPx_2)$. Clearly, $N_B(r) \neq \emptyset$.

Suffices that we prove that $N_B(r) \setminus \{y\} \neq \emptyset$. Indeed, if so, and $r' \in N_B(r) \setminus \{y\}$, then an $(x_4, \{y, r'\})$ -2-fan in B (which exists as B is nontrivial by (4.6.A)), a $y\ell$ -path in $H' - x_5$, where $\ell \in N_{H'-x_5}(x_5)$, and the edges $\{x_5r, rr', x_5\ell\}$ define a circuit satisfying (F.2).

Suppose then that $N_B(r) = \{y\}$. If so, then the following terms are satisfied:

- (i) $\{y, z, r\}$ is a triangle; indeed, $rz \notin E(P)$ implies that $zPr \subset N_G(y)$ and thus that $K_4^- \subset G$.
- (ii) $rx_3 \in E(G)$; implied by planarity and the assumption that $N_B(r) = \{y\}$.
- (iii) $zx_3 \notin E(G)$; or $K_4^- \subset G$ by (i) and (ii) above.
- (iv) $z \neq z'$; otherwise H' consists of the single edge x_5y and thus $\{z, y, r, x_5\}$ induce a K_4^- .
- (v) $N_G(x_3) \cap (zPx_1) = \emptyset$, meaning that $N_G(x_3) = x_1 \cup Y$ such that $Y \subseteq rPx_2$. To see this, we show that if there is a $u \in N_G(x_3) \cap (zPx_1)$, then $TK_5 \subseteq G$ which is constructed as follows:
 - (v.a) $\{r, x_5, x_2, x_3\}$ are the branch vertices of a TK_4 . Indeed, $\{r, x_5, x_2, x_3\}$ induce a C_4 and the paths rPx_2 and (x_5, x_1, x_3) complete this C_4 into a TK_4 .
 - (v.b) Let $F = \{P_1, P_2, P_3, P_4\}$ be the following $(y, \{r, x_5, x_2, x_3\})$ -4-fan: P_1 consists of the edges $\{yz, rz\}$. P_2 is a yx_5 -path in H' . Let P'_3 and P'_4 be the members of a 2-fan in B from y to $\{x_4, \ell\}$, where $\ell \in N_B(z') \setminus \{x_4\}$; such a vertex exists for otherwise planarity would imply that x_1 is at most 4-valent in G .
 - (v.c) Let P_3 and P_4 be obtained from P'_3 and P'_4 by adding the edge x_4x_2 to the member of $\{P'_3, P'_4\}$ that is a yx_4 -path. The other member of $\{P'_3, P'_4\}$ is a $y\ell$ -path. Add to it the edge $z'\ell$ and the subpath $z'Pu$ and the edge x_3u .

The union of F and the TK_4 constructed in (v.a) form a TK_5 .

We may now assume that $N_G(x_5) \cap (z'Px_1) = \emptyset$; otherwise, by a symmetrical argument to the one above we will have that $N_G(x_3) \cap (z'Px_2) = \emptyset$ contradicting (ii). Thus, $\{z', y, r\}$ is a 3-disconnector of $G - a$ separating x_3 from $H' - x_5$ which is nonempty as seen in (iv); contradiction. $\square$

Proof of (4.6.B.3). If $z = z'$, then H' consists of the single edge x_5y ; consequently, x_5 is adjacent to at least one of $(z'Px_1)$ or (zPx_2) contradicting (4.6.B.2). Next, if, say, $x_1 = z'$, then either $\{x_1, y, z, x_2\}$ is a 4-disconnector of G or $\{x_1, y, z, x_2, x_3\}$ is the boundary of a planar hammock since B is nontrivial, by (4.6.B.1). $\square$

Proof of (4.6.B.4). Suppose not. Then $\{x_1, z, y, z', x_2\}$ are the boundary of a 5-hammock of G of order 6 (otherwise we have a planar hammock of G); implying that B is trivial, contradicting (4.6.B.1). $\square$

Proof of (4.6.B.5). Suppose H' consist of the single edge x_5y . The segment $z'Pz$ consists of a the single edge zz' then; for vertices in $(z'Pz)$ can only have degree ≤ 4 since x_3 is a singleton P -bridge. Thus, $z, z' \in N_{G-x_5}(x_5)$, by (4.6.B.2); implying that $\{z, z', x_5\}$ is a triangle. Thus, neither of z, z' is adjacent to y ; or $K_4^- \subseteq G$. In addition, at least one of z, z' , say z , is not adjacent to x_3 ; or $K_4^- \subseteq G$. To be at least 5-valent, then, $|N_B(z)| \geq 2$. We then construct a $\{z, x_5, x_4\}$ -circuit satisfying (F.2) as follows. The union of the edges $\{x_5y, x_5z, z\ell\}$, where $\ell \in N_B(z) \setminus \{y\}$, and an $(x_4, \{y, \ell\})$ -2-fan in B form such a circuit. $\square$

Proof of (4.6.B.6). Suppose not. The set $\{x_5, y, w, w'\}$ has size ≤ 4 . Since this set is not a 4-disconnector, H' consists of the single edge x_5y ; contradiction to (4.6.B.5). $\square$

Proof of (4.6.B.7). If such a path does not exist, then $\{x_4, y\}$ is a 2-disconnector of B

separating h'_2 from b . That is, $B = B_1 \cup B_2$ such that $G[\{x_4, y\}] = B_1 \cap B_2$, $h'_2 \in V(B_1)$ and $b \in V(B_2)$. Since $|\{x_1, x_2, z, z'\}| = 4$, by (4.6.B.3), $N_{B_i}(x_i) \setminus \{y, x_4\} \neq \emptyset$ for $i = 1, 2$. As neither of $\{x_1, z', y, x_4\}$, $\{x_2, z, y, x_4\}$ is a 4-disconnector of G , it follows that x_3 is adjacent to both $(x_1 P z')$ and $(x_2 P z)$. Since we assume that neither of $\{x_3, x_1, z', y, x_4\}$, $\{x_3, x_2, z, y, x_4\}$ form the boundary of a planar hammock, each of these sets is the boundary of a 5-hammock of G of order 6. This then implies that x_3 is adjacent to B ; contradiction to the premise of the DISCONNECTED CASE. $\square$

■

References.

- [1] R. Diestel, *Graph Theory*, third edition, Springer, 2005.
- [2] J. Ma and X. Yu, Independent paths and K_5 -subdivisions, J. Combinatorial Theory B (to appear).
- [3] J. Ma and X. Yu, K_5 -subdivisions in graphs containing K_4^- , submitted manuscript.
- [4] W. Mader, $3n - 5$ edges do force a subdivision of K_5 , *Combinatorica*, **18** (4) (1998) 569-595.
- [5] D. M. Mesner and M. E. Watkins, Cycles and connectivity in graphs. *Can. J. Math.*, **19** (1967) 1319 – 1328.
- [6] B. Mohar and C. Thomassen, *Graphs on Surfaces*, 2001, The Johns Hopkins University Press Baltimore and London.
- [7] P. D. Seymour, Disjoint paths in graphs, *Discrete Math.*, **29** (1980) 293 – 309.
- [8] C. Thomassen, 2-linked graphs, *European J. Combin.*, **1** (1980) 371 – 378.
- [9] X. Yu, Subdivisions in planar graphs, *J. Combin. Theory Ser. B*, **72** (1998) 10 – 52.